

ENDOSULFAN

Fact sheet
&
Answers to common questions



Thanal

L-14 Jawahar Nagar, Kowdiar, Thiruvananthapuram - 695 003
Kerala, India

Thanal is a voluntary public interest research, advocacy and action group working on matters of Environmental Health and Justice. It has specific areas of work in Zero Waste, Pesticides and Health, Natural History, Earth Education, and Toxics in the environment. Issues related to industrial pollution, pesticide contamination, wrong and inappropriate technologies are a major area of the work.

Thanal is a participating organization in IPEN and co-chair of the Pesticide Working Group and officiates as its International Secretariat.

This fact sheet was developed after four years of interaction with experts and scientists all over the world. Harikrishnan V R, Researcher and Usha S, Coordinator of the Pesticide and Agriculture Programme of Thanal compiled the fact sheet.

The organisation is indebted to IPEN-Pesticide Working Group, IPEN-Community Monitoring Working Group, Pesticide Action Networks of Asia-Pacific, United Kingdom and North America. Thanal would have much wished to name and acknowledge all the people who have helped technically and personally, had the list not been too big.

Nevertheless, Thanal wishes to acknowledge the single most driving spirit behind such a compilation - the people in the villages of Kasaragod, whose suffering is probably incomparable and their fight against the pesticide Endosulfan the only fight of this kind for survival.

Endosulfan - A Fact sheet and Answers to common questions
Harikrishnan V R and Usha S
January 2004

Thanal

L-14 Jawahar Nagar, Kowdiar,
Thiruvananthapuram - 695 003
Kerala, India.
Telephone: ++91-471-2727150
Email : thanal@vsnl.com
Website : www.thanal.org

ENDOSULFAN

A Fact Sheet and Answers to Common Questions

Introduction

Endosulfan is a pesticide belonging to the organochlorine group of pesticides, under the Cyclo diene subgroup. Introduced in the 1950's, it emerged as a leading chemical used against a broad spectrum of insects and mites in agriculture and allied sectors. It acts as contact and stomach poison and has a slight fumigant action¹. It is used in vegetables, fruits, paddy, cotton, cashew, tea, coffee, tobacco and timber crops. It is also used as a wood preservative and to control tse-tse flies and termites⁵. It is not recommended for household use. Intentional misuse of endosulfan for killing fish^{2,57} and snails⁷¹ has also been reported.


Endosulfan was introduced at a time when environmental awareness and knowledge about the environmental fate and toxicology of such chemicals were low and not mandatory as per national laws. But now it is being detected as an important cause of pesticide poisoning in many countries.

Chemical Name

6,7,8,9,10,10- hexa chloro- 1,5,5a,6,9,9a- hexahydro – 6,9- methano- 2,4,3- benzodioxathiepine-3-oxide.

Chemical Formula $C_9H_6Cl_6O_3S$

Chemical Structure:-



In pure form endosulfan exists as colourless crystals. But the technical product is brownish crystals with slight odour of sulphur dioxide³. Technically endosulfan is a mixture of two isomers - alpha-endosulfan and beta-endosulfan in the ratio 7:3. Technical grade endosulfan contains 94% alpha-endosulfan and beta-endosulfan and other related compounds like endosulfan alcohol, endosulfan ether and endosulfan sulfate. Endosulfan is insoluble in water, but it dissolves in xylene, chloroform, kerosene and most organic solvents and is a noncombustible solid. It is mixable with most fungicides and compatible with most pesticides⁴.

Toxicity Classification

The U S Environmental Protection Agency (EPA) classifies endosulfan as Category Ib – Highly Hazardous. The European Union also rates it Highly Hazardous. World Health Organisation (WHO) classifies endosulfan in Category II - Moderately Hazardous. The classification of WHO was found to be inappropriate considering the classification followed in countries all over the world and the available toxicity information. It has been alleged that the classification is based mainly on LD50 value for acute toxicity generated by the producer company¹³. The Industrial Toxicological Research Centre (ITRC) in India the nodal centre for the Regional Based Assessment of Persistent Toxic Substances (PTS) for the Indian Ocean region by the UNEP-GEF classifies endosulfan as Extremely Hazardous⁴.

Q. What do international treaties say about endosulfan ?

Endosulfan is widely considered to be a Persistent Organic Pollutant (POP) but was not included in the initial list targeted for phase out under the Stockholm Convention. Endosulfan was in the initial list of POPs being considered for world-wide elimination at the first meeting of experts in Vancouver, Canada (1994) jointly convened by governments of Canada and Philippines but was later removed from the list¹³. Endosulfan is recognized as a Persistent Toxic Substance (PTS) by the UNEP.² It is also considered as a potential POPs candidate.

Q. What is the regulatory status of endosulfan in world nations ?

Endosulfan has been in world-wide use since its introduction in the 1950's. It was considered a safer alternative to other organochlorine pesticides in many countries all over the world since the 1970's. But in the last two decades many countries have recognized the hazards of wide application of this pesticide and has either stopped its production and/or banned or restricted its use.

Endosulfan is banned in countries like Singapore, Belize, Tonga, Syria, Germany, Sweden, Philippines, Netherlands, St. Lucia, Columbia, Cambodia, Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, UAE, Sri Lanka and Pakistan^{2,73}. Restricted use is allowed in Australia, Bangladesh, Indonesia, Iran, Japan, Korea, Khazakhstan, Lithuania,

Thailand, Taiwan, Denmark, Serbia & Montenegro, Norway, Finland, Russia, Venezuela, Dominican Republic, Honduras, Panama, Iceland, Canada, US and UK^{2,56,73,80}.

It is one among the twenty one priority compounds identified by the UNEP-GEF in the Regional Based Assessment of Persistent Toxic Substances (PTS), 2002. These reports have taken into account the magnitude of use, environmental levels and human and ecological effects of these compounds². In the Indian Ocean Region endosulfan is banned in 8 nations. India is one of the major Indian Ocean rim nations, which has imposed no ban or restrictions on endosulfan. A ban on endosulfan exists in the South Indian state of Kerala (imposed through a Court Order), which came as a result of a public pressure following the poisoning of many villages due to aerial spraying of the chemical⁸⁴. Colombia and Cambodia are two countries where endosulfan is banned from use recently⁸⁰.

In the Reports of the Regional Based Assessment of PTS of the UNEP-GEF, endosulfan has been rated depending on the level of concern in the respective region countries. They are as follows-

- **Indian Ocean region** – “Regional Concern”²
- **North American region**– Treated as “Regionally specific PTS”, receiving great attention along with HCH, Chlordane, PCB, PAHs⁷⁴.
- **Mediterranean region** – “Local Concern”⁷⁰.
- **Sub Saharan Africa** – Ranked as the PTS of highest concern after DDT⁶⁹.
- **Eastern and Western South America** – “Potentially relevant PTS of emerging concern”⁷².
- **European region** – “Proposed Possible Priority Hazardous Substance”⁷⁵.
- **South East Asia and Southern Pacific region** – “Regional Concern”. It is identified as a major PTS which has a continuing effect on the natural ecosystem in the region and long term effect on the structure of aquatic ecosystem⁷¹.
- **Central America and the Caribbean** – Considered as one of most important PTS of emerging concern⁷³.

Q. How much endosulfan is produced, used and stockpiled ? and Where ?

Practically very little information is available of the volumes of production of endosulfan. WHO estimated that the worldwide production of

endosulfan was 10,000 MT in 1984⁸¹. Current estimates of worldwide production or domestic formulations could not be located⁸¹. Many countries that produced endosulfan in the 1970's and 80's do not produce it any more. For instance endosulfan has not been produced in the United States since 1982, but it is still used in chemical formulations.⁸¹ India is one of the major producers of endosulfan. Since 1996-97 it produces an average of 8206 MTPA totaling 41033 MT between 1995-2000². India exported 12180 MT during this period and consumed on an average 3599 MTPA. The UNEP-GEF reports on PTS has identified some of the producing and importing countries but there is a large data gap. No information regarding stockpiles of endosulfan could be located.

Q. How might I be exposed to endosulfan ?

- By breathing air near where it has been sprayed.
- Drinking water contaminated directly or through run-offs.
- Being in contact with contaminated soil.
- Eating contaminated food.
- Smoking cigarettes made from endosulfan contaminated tobacco.
- Working at endosulfan production centers.
- Direct ingestion.

Q. How does endosulfan affect human beings ?

Worldwide use of endosulfan increased with the ban/restriction in use of the more persistent organochlorine pesticides like DDT and Endrin. Much research has revealed off-late that this chemical is also as problematic to life forms, the effects being comparable to those exhibited by its predecessors in the organochlorine class.

Endosulfan is highly toxic and can be fatal if inhaled, swallowed or absorbed through the skin. A number of acute poisoning cases have been reported. Ingestion or breathing high levels of endosulfan may lead to convulsion and death. Endosulfan directly affects the central nervous system^{24,28}. It is absorbed through skin and eye irritation may also result. Symptoms of poisoning include hyper activity, excitement, dyspnea (breathing difficulty), apnea (stoppage of breathing), salivation, loss of consciousness, diarrhea, anemia, nausea, vomiting, insomnia, blurred vision, cyanosis (bluish discoloration of skin due to want of oxygen), foaming at the mouth, tremor, dry mouth, lack of appetite, irritability, head ache, decreased respiration, loss of memory, haematuria, albuminuria, confusion, dizziness, imbalance and lack of coordination^{4,5,7,30}. Persons suffering from asthmatic and convulsive

disorders are at high risk. Persons on protein deficient diet also possess high risk^{24,30}.

Autopsy examination of a intentional ingestion (suicide) case has revealed damage to liver, lung and brain^{8,8a}. Endosulfan is a proven endocrine disruptor^{6,9}. It has potential to induce hypo thyroidism⁶⁶. Long term health effects are not properly studied, experimented or documented world wide. Acute toxicological data is however adequately available. Acute oral toxicity is higher than dermal toxicity³⁰. Absorption rate and toxicity is found to increase in the presence of solvents like alcohol and aromatic solvents^{12,24,34,61}.

Endosulfan exhibits estrogenic properties^{9,43}, comparable to that of DDT⁹. It competes for estradiol for binding to estrogen receptors, thereby inhibiting hormonal function. It induces proliferation of human breast estrogen sensitive MCF7 cells⁴⁹, (invitro) thereby increasing breast cancer risk. It harms the reproductive system by affecting semen quality, sperm count, spermatogonial cells, sperm morphology and other defects in male sex hormones¹¹. It is found to inhibit testicular androgen biosynthesis in lab animal experiments³⁰ and exhibits significant risk in renal and testicular damage. It may have adverse effects on central nervous system by inhibiting brain acetyl cholinesterase¹⁶, causing uncontrolled discharge of acetyl choline. Endosulfan ingestion is known to affect the kidneys and liver⁵⁰. It inhibits leucocyte and macrophage migration (this is the inhibition of the natural immune system by disrupting anti-body protection) causing adverse effects on humoral and cell-mediated immune system³⁰.

Endosulfan is found to damage human red blood cells (RBC) at concentrations of 1ppb-1ppm¹⁰. Both alpha-endosulfan and beta- endosulfan are genotoxic to HepG2 cells⁸. Endosulfan is hepatotoxic⁶, mutagenic, clastogenic and induce effects on cell cycle kinetics¹³. Endosulfan is known to cause chromosomal aberrations in hamster and mouse and sex linked mutations in *Drosophila*^{13,30,58}. Endosulfan has caused mutations in bacterial and yeast cell. It is also known to cause mutations in mammals¹³. It may induce mutations in human, if exposure is great. It is also a potential tumor promoter⁶⁷.

Many studies related to its acute and chronic toxicity in laboratory animals are available. Endosulfan is highly toxic to rats and mice^{13, 30}. Some studies suggests its teratogenic²⁸ and carcinogenic properties²¹ on rats and mice. It directly affects the central nervous system, causes liver and kidney (chronic glomerulonephrosis) damage⁶ in rats and mice. It also impairs the reproductive system of rats³⁹. Behavioural and neurological changes have also been noticed³⁰. Thyroid follicular damage in mouse has been reported^{30,66}. Endosulfan is known to damage the endocrine system, nervous system,

circulatory, reproductive, respiratory and excretory systems and developing foetus.^{6,7,14,15,16,21,30,37}

The National Institute of Occupational Health (India) have linked the higher prevalence of neurobehavioral disorders, congenital malformations in female children and abnormalities related to male reproductive systems to the continuous exposure to endosulfan spray. The study was conducted among children in one of the villages in Kasaragod District (in the South Indian State of Kerala) where endosulfan was aerially sprayed⁶⁴. Endosulfan is implicated in the occurrence of adverse health effects particularly in rural communities in South East Asia, Southern Pacific and Sub-Saharan Africa.^{69,71}

Q. Can endosulfan cause cancer ?

No accurate data related to the carcinogenicity of endosulfan in human is available. But its carcinogenicity has been proven in male and female Osborne-Mendel rats²¹. Endosulfan induced malignant neoplasms at all sites in male and female rats and endocrine organs in male rats²¹. Both sexes developed lymphosarcomas and female rats had neoplasms of the reproductive system²¹. Endosulfan is also carcinogenic for the liver of female mice²¹. But from field level reports, endosulfan can be rightly suspected for having carcinogenic properties in human beings, especially in cases of chronic exposure. In some reports it is referred to as having possible carcinogenic effects, effects in human immune and reproductive system⁶⁸. Studies have also proven that it induces proliferation of human breast estrogen sensitive MCF7 cells in vitro^{9,49} which may lead to greater breast cancer risk.

Q. What happens to endosulfan in the environment ?

The fate of endosulfan released in the environment is different for the two isomers and also depends on the medium it gets deposited.

Beta- endosulfan is more persistent than its alpha- isomer. Endosulfan sulphate is the main degradation product of both isomers, which is equally toxic^{37,66} and is itself more persistent in the environment than its parent compounds^{5,30}. Endosulfan can be broken down by photolysis, hydrolysis and bio degradation. Endosulfan diol, endosulfan lactone, endosulfan ether etc. are some of the other byproducts⁵. Although the isomers are fairly resistant to photolysis, the break down products are susceptible. On plant surfaces endosulfan rapidly degrades to metabolites³⁰. Endosulfan is fairly immobile in soil and is highly persistent^{13,33}. Major products in soil are endosulfan diol, endosulfan sulfate and endosulfan lactone. Endosulfan sulfate production increases with an increase in

temperature³⁰. Endosulfan will persist longer under more acidic conditions^{5,24}. It persists longer under submerged conditions⁸⁶. The half life of endosulfan varies from 60 days (alpha- endosulfan) to 800 days (beta-endosulfan)¹³. It enters air by volatilization from plant and water surfaces^{13,33}. Contamination by drift and particle transport also occur³³. Ultra low volume (ULV)- application can drift several kilometers from point of application³⁰. It has been detected from far off place like Arctic air, snow-water and lake waters, rainfall and snow samples in Californian mountains and remote European mountain lakes^{5,30,33,68,75}. Beta-endosulfan is more stable in air. Endosulfan has been shown to be released from wood preservatives into room atmosphere over an one year period of observation⁵. In water endosulfan has a half life of 35 to 150 days¹³. It does not easily dissolve in water and may accumulate in bodies of fish and other aquatic organisms^{30,69}. The break down products in water are endosulfan-diol and endosulfan sulfate. Endosulfan does not reach down much to ground water³⁰, but has been proven to run off after spraying. But it has been detected from ground water at deep soil layers in concentration ranging from 0.009- 0.053 micro gram per litre upto 20 days after last spraying¹³. USEPA recommends that the levels of endosulfan in rivers, lakes and streams should not be more than 74 ppb²⁸. But this limit is 15 times more than the concentration required to cause reproductive damage in red spotted newt⁵⁴. Several studies of bioaccumulation have been conducted around the world and residues were found in aquatic organisms. USEPA considers endosulfan as having a high bioaccumulation potential in fish¹⁶, but not much evidence of bioaccumulation in higher trophic levels is available.

Q. Is there any evidence of endosulfan contamination in the environment, food and human beings ?

Reports of endosulfan residue in food, soil, air, body tissues etc are available from all parts of the globe.

Residues were detected from air, water (surface and ground water)^{2,59} and soil in India², water and sediments in Ghana⁴⁵, marine water and sediments in India², lagoons in Spain, coastal, estuarine and river sediments in Israel (high concentrations in Lake Kinneret)⁷⁰, water in Benin, Malawi, Nigeria and South Africa (alarming levels in river water 684-4843 ng/L), soil in Benin, Nigeria, Sudan and Zambia, sediments in Benin and Nigeria, vegetation in Madagascar, Zambia and Ghana, Water from remote mountain lakes in Europe (the Alps, Caledonian and the Pyrenees)⁷⁵ and river and sea water in South East Asia⁷¹ etc. were found to be polluted with endosulfan. In Malaysia river sediments were found to have high levels (434micro grams/litre)

of endosulfan⁷¹. Reports from Central America shows air in Costa Rica, coastal waters, estuaries, well, surface and marine waters and sediments in Honduras and Jamaica contaminated with endosulfan residues⁷³. Endosulfan is one of the most frequently reported PTS in surface and ground water in Central America and the Carribean⁷³. In Guatemala, it is found more frequently and at higher levels in river water, well water and spring water⁷³. It has been detected in municipal water system in Colombia⁷³.

High concentrations of alpha and beta endosulfan isomers and endosulfan sulphate have been detected in tree bark samples through out the world, particularly in India and the Pacific Rim³⁰.

Endosulfan has been detected from food samples²⁰ around the world- in Australian beef at 0.36 mg/kg³² (2 times the Australian limit and 4 times the international limit), in cows milk from tobacco farming areas in USA and food samples in USA and Canada⁵. Residues were detected in tomatoes from Brazil²⁰, Spanish pepper samples from Finland, fish from India², Kenya and Nigeria⁶⁹, food and vegetables in Croatia⁷⁰ and vegetables in Cyprus⁷⁰. High levels of residues were found in red pepper and egg plants in Catania(Italy)⁷⁰. High residues have also been found in diary food, meat, fish, chicken, and vegetables in Eastern and Western South America⁷², cows milk in Brazil⁷⁹, vegetable diet in Kuwait, vegetables, vegetable oil and seeds from India and animal samples from slaughter houses in India². Endosulfan residues were detected from cows milk in Colombia far exceeding the WHO and FAO reference levels⁷³. Fish from Kenya and Nigeria were heavily contaminated with endosulfan⁶⁹. The European Union has banned import of fish from Tanzania, Uganda and Kenya due to high levels of endosulfan residues⁸⁰. Residues were also detected from animals samples in Benin, Nigeria, Cote d' Ivoire, Madagascar, South Africa and Kenya⁶⁹. Presence of endosulfan in Lake trout in North America suggests a wide dispersal from areas of use to isolated lakes⁷⁴.

Endosulfan has also been detected from human tissues. It has been detected from cord blood samples obtained at the time of delivery⁴⁶, human sera^{42,55} and human milk samples obtained from healthy lactating women in Spain⁴⁴. Human breast milk from Egypt^{30,41}, Colombia and Nicaragua⁷³ and from cotton pickers in Pakistan² were found to be contaminated. High levels of endosulfan were detected in human breast milk in Sub Saharan Africa⁶⁹. Residues were also detected in fat samples from children living nearby farms in Spain²². Blood, human milk and urine samples in Croatia were also found to be contaminated. Alarmingly high levels of endosulfan residues were observed in human blood and milk in a study in Kasaragod in Kerala, India^{26,26a,27,31,64}.

Q. How does endosulfan affect wildlife and domestic animals?

There are many reports regarding the toxicity of endosulfan on wildlife. National Wildlife Federation (USA) states that endosulfan is extremely toxic to wildlife and acutely toxic to bees¹³. It is acutely toxic to birds - mallard ducks²⁴, quails and pheasants³⁰.

Endosulfan is highly toxic to aquatic organisms even at recommended levels of application^{13,71}. It is particularly toxic to fishes^{5,18,24} – massive fish kills are reported from many places¹³. It also causes endocrine problems, reduction of protein in tissues and other health effects. The high toxicity to fish species is evidenced through studies on *Gambusia affinis*¹⁸, Rainbow trout, channel catfish, bluegill sunfish, minnow^{24,30,36}. It is known to impair the pheromonal systems leading to disrupted male choice and lowered mating success in female red spotted newts⁵⁴. It is known to affect the germ cell population of zebrafish embryos⁴⁷. It exhibits anti-estrogenic effects in fresh water catfish, reproductive problems in female teleostfish and opercooidfish³⁰. It also bioconcentrates in aquatic organisms³³. Reports from South East Asia and Southern Pacific proves that endosulfan has detrimental effects on aquatic biota⁷¹. Reports prove its high toxicity to frogs, toads, annelids, aquatic insects (damselflies, midges, beetles etc), crustaceans (crab, shrimps, prawns etc), fishes and molluscs⁷⁸. Endosulfan affects the hatching rate and larval survival of common Indian Toad⁸².

Besides being toxic to crab larvae²⁹, it has been reported to disrupt the moulting of crabs and aquatic invertebrates^{76,76a}. It is known to function as an endocrine disruptor in American alligators⁶⁵. It has been detected from the nonviable eggs of Morelets crocodile¹⁹. It is also known to be affecting the egg laying of grass shrimps⁵². It has been detected as one of the predominant organochlorine pesticides in the silverside fish in Argentina⁵³. It has been related to the drastic population decline of anuran amphibians in Western USA in over the last 10 to 15 years⁴⁸. Residues have been obtained from the liver of Eastern Box turtle⁴⁰. It is found to accumulate 600 times the water concentration in mussels²⁴.

It is toxic to non target organisms¹⁶, like predators of several pests⁵¹. Endosulfan is highly toxic to soil micro arthropods¹⁷, micro organisms, phytoplanktons, soil algae, actinomycetes, bacterial colonies etc.^{13,30}. It is also toxic to mammals like rabbits^{13,30}. The disappearance of cats, frogs, bees, fresh water fishes etc. were reported from Kasaragod District in South India, where endosulfan was aerially sprayed²⁷.

Endosulfan is also highly poisonous to cats (LD50- 2mg/kg) and dogs (LD50-76.7mg/kg). Reversible blindness and lack of muscle coordination has been reported in sheep and cattle grazing on endosulfan contaminated grass²⁴. Farmers from

endosulfan sprayed area report of acute effects to cattle at time of spray and after, and leading to fatalities²⁷.

Endosulfan exhibits phytotoxic properties⁷. Toxic effects on plants like root growth inhibition, stunting, burning of tips and margins of leaves and affected root permeability have been reported¹³. It is a prominent contaminant in vascular plants and lichen even in remote areas like the Arctic⁶⁸.

Q. Is there any reports of endosulfan poisoning from around the world ?

Cases of endosulfan poisoning have been reported from many parts of the world. Accidental and intentional exposure leading to human fatalities and environmental tragedies has occurred. The following are some of the major cases of poisoning.

A wide spread fish kill was observed in 1969, when 30 kilograms of endosulfan was discharged into the Rhine river in Federal Republic of Germany^{5,30}. In 1975, an accidental spill of endosulfan caused a major fish kill in North Brook a tributary of the Dunk River in Eastern Prince County of Prince Edward Island. The Brook trout population was reduced from 2227-4147 to 45-246³⁰. In the Philippines, endosulfan accounted for the largest number of deaths due to pesticide poisoning reported in 1991¹³. In 1995, run-off from cotton fields contaminated with endosulfan resulted in the death of more than 24,000 fish along a 25-kilometer stretch of a river in Alabama¹³. Investigations showed that the pesticide had been sprayed according to label instructions. In Sudan in 1988, endosulfan barrels washed in irrigation canals caused fish death. In addition three people died after drinking water from the canal¹³. In Northern India, 18 cases of endosulfan poisoning have been reported in 1995-97 by accidental over-exposure during spray²³. In 2000, a case of 44 individuals who consumed food accidentally contaminated by endosulfan was reported in rural India. One individual died⁴⁴. In Sulawesi, Indonesia, 32 cases of poisoning due to endosulfan have been reported from 1990 to 1993⁵⁶. In Columbia, 155 cases of poisoning (in 1994) and 60 cases (1993) of poisoning due to endosulfan were reported⁵⁶. The misuse of endosulfan to kill snails has resulted in the largest number of poisoning cases with fatalities in Philippines in 1996⁷⁵. Many cases of poisoning-death in Guatemala, Costa Rica etc. have also been reported⁷³. In Feb 2002, two South African boys living near Ntabamhlophe, Kwa-Zulu Natal died following exposure to endosulfan. A police officer and 3 journalists were hospitalized after visiting the place several hours later⁸⁰. The worst of all the cases so far reported are from three nations- Cuba, Benin and India.

Pesticide Poisoning in Cuba : Endosulfan was responsible for the deaths of 15 people in the western province of Matanzas, Cuba in February 1999. A total of 63 people became ill after consuming food contaminated with endosulfan according to Cuban authorities⁸³.

The Benin Tragedy:- In Borgou province in Benin, Endosulfan poisoning caused many deaths during 1999 - 2000 cotton season. Official records state that atleast 37 deaths occurred and 36 were taken seriously ill. In the same region in 1999 a boy died after eating corn contaminated with endosulfan²⁵.

The Kasaragod Tragedy, South India:- People in 15 villages in Kasaragod in the South Indian State of Kerala were subjected to continuous exposure to endosulfan which was aerielly sprayed three times every year for 24 years. Congenital Birth defects, reproductive health problems, Cancers, loss of immunity, neurological and mental diseases were reported among the villagers. Following a public outcry a number of health based scientific studies confirmed that the health problems were directly linked to the exposure to endosulfan^{27,60,64}.

Q. What are the alternatives to endosulfan?

While there are many non-chemical alternative practices, the choice of one primarily depends on the crop and the pest for which it has been used.

One of the important crops on which endosulfan is used is cotton. Endosulfan is also used on crops like vegetables, cashew, coffee, tea etc.

There are numerous documented benefits from the adoption of Integrated Pest Management (IPM) programme for cotton from different cotton growing areas all over the world. The focus of IPM is on cultural and biological control of pests rather than depending on pesticides. A comparison in Frio county, Texas show a savings of between 131- 300 percent for alternative cotton pest management⁶². Organic farming of cotton is also gaining popularity among cotton farmers in the Asian region⁶².

Similarly in vegetable cultivation a lot of field trials has been done on alternative pest control methods focusing on herbal pesticides. In the Asian region farmers have developed their own combinations and methods of pest control using chillies, garlic, asafoetida, cow urine and many other plant materials. FAO supported IPM programme in vegetables is well documented.

In India endosulfan was recommended for controlling tea mosquito bug in Cashew and Tea until 2000. Following the Kasaragod Tragedy, the National Research Centre for Cashew has withdrawn its recommendation for endosulfan use in cashew.

Many cashew farmers have tried organic method and application of neem oil and pongamia leaf extract is found successful by some South Indian farmers. Some farmers have even tried 'do nothing' farming and the results are encouraging. Alternative pesticides and organic farming are encouraged in tea plantations and the use of endosulfan is being eliminated. There are tea plantations in South India which has adopted bio-dynamic farming and have completely eliminated the use of chemical pesticides including endosulfan.

In coffee, to control coffee berry borer, Integrated Pest Management is adopted instead of endosulfan⁶³. This includes strict phytosanitary and cultural measures supplemented with bio control measures and mass trapping of the insects.

Though alternatives to endosulfan are available, support for such practices are very low. If Governments and research institutions can support such work, use of endosulfan can be totally eliminated in agriculture and other sectors.

Q. What can I do to eliminate the use of endosulfan?

- Look out for the alternative non-chemical pest control which is available in your region and other parts of the world and thus eliminate the use of endosulfan.
- Pressure your government to ratify the Stockholm Convention if it has not already done so (see <http://www.pops.int> for a current list) and press for addition of endosulfan to the Stockholm Convention List.
- Pressure your Government to ban the production and use of endosulfan, if it has not already done so.
- Join one of the local and international campaigns to ban endosulfan.
- Adopt "Land and Food without Poisons" as a goal for survival and adopt organic, ecological or natural agriculture in your farm. Share this message among fellow farmers.

Trade names and formulations for endosulfan:-

Endosulfan is available in the market in a number of trade names. It is available as formulations of Emulsifiable Concentrate (EC), Wettable Powder (WP), Ultra low Volume Liquid (ULV), Granules (G), Dust(D) and Smoke Tablets.

The following list contains some of its trade names. The list is not complete and some names may now be obsolete.

Bangladesh – Thiodan

Brunei – Thiodan, Fezdion

Chile – Parmazol E, Flavylon, Galgofan, Thiodan, Thionex, Thionyl y methofan

India – Agrosulphan, Agiro Sulphan, Banej Sulphan, Cilo Sulphan, Endo Sulphan, E- Sulfan, Endo Chithin, Endocid, Endonit, Endomil, Endosol, Endostar, Endosun, Endotaf, Endostan, Endocing, Endocide, Endosulpher, Gaydan, Gilnore Endorifan, Hexa-sulfan, Hildan, Hockey Endosulfan, Hy-sulfan, Kemu Sulfan, Hilexute-Sulfan, Krushi Endosulfan, Lusu Sulfan, Marvel-Micosulfan, Mico Thansulfan, Pary Sulfan, Pesticel, Remisfan, Sico sulfan, Solesulfan, Sujadin, Sulfan, Tej Sulfan, Thiodon, Thiokill, Thionel, Thionex, Thioton, Veg-fru Thiotox, Veg-fru Thiotex, Vika sulfan.

Indonesia – Thiodan, Fanodan, Dekasulfan.

Korea – Malix, ThioliX.

Pakistan – Siagon, Thiodan, Thioluxan.

Philippines – Atlas Endosulfan, Endosulfan, Contra, Endox, Thiodan.

Sri Lanka – Thiodan, Thionex, Endomack, Endocel, Bours Endosulfan, Harcos Harcosan, Red Star Anglo-sulfan.

Thailand – Thiodan, A. B. Fan, Agrodan, Agridan, Bensodan, Bensocarb, Beosit, Brook, Clement, Dew Dan, Dior 35, Dori, Dumpersan, E C Sulfan, Egodan, Endan, Endodan, Endosulfan, Endrew, Endye, Endyne, Etonic, Exxo-Z, Famcodan, Fortune, Freedan, Gardner, Gycin, Hor Mush, Hydrodan, J-teedan, Jack Dum, Kasidan, L P dan, Lordjim, Malix, Manyoo, Metrodan, Nayam, Newcodan, Nockdyne, Ox Xa, Patodan, Pestdye, Pro-d-dan, Sandan, Shevanex, Simadan, Sonydan, Summer, Tanadan, Teophos, Thanyacarb, Thimul, Thiofor, Torpidan, Urofen, Wephos, Zumic.

Other Names: Chlorthiepin, CycloDan, Endox, Thifor, Thiomul, Thionate.

References

1. K.K. Nayar, T.N Ananta Krishnan and David B.V. General and Applied Entomology. Tata McGraw Hill Publishing Co Ltd, New Delhi, India.

2. Anon (Dec 2002).
Regional Based Assessment of Persistent Toxic Substances- Indian Ocean Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility.

3. Anon (June 1997).
NIOSH Pocket Guide to Chemical Hazards. US Dept of Health and Human Services- National Institute for Occupational Safety and Health.

4. Anon (1989).
Toxicity Data Hand Book. Vol:III, Pesticide-A. Industrial Toxicology Research Centre(Council of Scientific and Industrial Research) Lucknow, India.

5. Anon (1984).
Environment Health Criteria 40- Endosulfan. IPCS (International Programme on Chemical Safety) – WHO Geneva.

6. Anon (22nd Dec 1997).
National Registration Authority for Agriculture and Veterinary Chemicals, Australia. Existing Chemical Review Programme – NRA Review of endosulfan – Draft.

7. Anon (Issue 2 May, 2000)
Hazardous Substances Data Book (HSDB)- US National Library of Media – Canadian Centre for Occupational Health and Safety.

8. Yuquan Lu, Kane Hiza MoriMoto, Tatsuya Takeshita, Toru Takeuchi, Takeshi Saito.
Genotoxic effects of A-Endosulfan & B-Endosulfan on Human HEPG2 cells – Environmental Health Perspectives ; JUNE 2000 ; 108-(6), 559-561

8a. Boerehoom F T, Van Dijk A, Van Zoonan P, Meulenbelt J (1998)
Nonaccidental Endosulfan Intoxication, A case report with toxicokinetic Calculations and Tissue concentration. J Toxicol. Clin Toxicol , 36: 345-352.

9. Soto AM; Chung K L, Sonnen Schein C(1994).
The Pesticides Endosulfan, Toxaphene and Dieldrin have estrogenic effect on human estrogen – sensitive cells. Environmental Health Perspectives – Vol 102, Iss 4; P 380-3.

10. Daniel C S, Agarwal S, Agarwal SS(1986)
“Human Red blood cell membrane damage by Endosulfan” Toxicol. Lett; Vol.32, ISS !-2, P 113-8. General and Applied Entomology. Tata McGraw Hill Publishing Co Ltd, New Delhi, India

11. Pandey N, Gundevia F, Prem A S, Ray P K. (1990)
Studies on the Genotoxicity of Endosulfan, an organochlorine insecticide in mammalian germ cells. Mutant Res; Vol. 242; ISS 1, P 1-7.

12. Fransis A Gunth (1968) (Edit)
Residue Reviews, Vol. 22.

13. Romeo F. Quijano, MD (Oct/Dec 2000).
Risk Assessment in a third world reality: An Endosulfan case History. International Journal of Occupational and Environment Health. Vol. 6, No. 4.

14. Paul V, Balasubrahmaniam E, Jayakumar A R, Kazi M. (1995)
A sex related difference in the neuro behavioural and hepatic effects following * chronic endosulfan treatment in rats. Eur. J. Pharmacol; Vol.293, ISS 4, P 355-60

15. Sinha N, Narayan R, Sankar R, Saxena D K. (1995).
Endosulfan induced bio chemical changes in testis of rats. *Vet. Human Toxicol*; Vol. 37, ISS 6, P 547-9.
16. Naqvi. S M, Vaishnavi C (1993)
Bio Accumulative Potential and toxicity of Endosulfan insecticide to non target animals. *Comp. Biochem. Physiol. C*; Vol 105, Iss 3, P 347-61.
17. Joy V C, Chakravarthy P P (1991)
Impact of Insecticide on Non Target Micro Arthropod Fauna in Agricultural Soil. *Ecotoxicol. Environ. Saf.* ; Vol. 22, ISS 1, P 8-16.
18. Karim AA, Haridi AA, el Rayah E A (1985)
The environmental impacts of four insecticides on non target organisms in the Gezira Irrigation Scheme canals of Sudan. *J. Trop. Medic. Hyg*: Vol 88, ISS 2, P 161-8.
19. Wu TH, Rainwater TR, Platt SG, Mc Murry ST, Anderson TA (March 2000)
Organo chlorine contaminants in Morelet's crocodile (*Crocodylus moreletti*) eggs from Belize. *Chemosphere*; 40(6); 671-8
20. Araujo A C, Telks D L, Gorni R, Lima L L (June 1999)
Endosulfan residues in Brazilian Tomatos and their impact on public health and the environment. *Bulletin Environ. Contam. Toxicol*; 62(6): 671-6.
21. Reuber MD , (Aug 1981)
The role of toxicity in the carcinogenicity of Endosulfan . *Sci. Total Environ.*; 20(1); 23-47.
22. Olea- Serrano F, Olea N, Lardelli-Claret P, Rivas A, Barba-Navarro A. Jan-Mar (1999).
Inadvertant exposure to xenoestrogens in children. *Toxicol Ind. Health*, 15(1-2); 151-8.
23. Chugh.S.N, Dhawan R, Agarwal N, Mohajan S K, (Sept 1998).
Endosulfan poisoning in north India, a report of 18 cases. *International Journal Clin. Pharmacol Ther*; 36(9); 474-7
24. Anon(1996)
Pesticide information Profile- Endosulfan (Revised June 1996)EXTONET- Extention Toxicology Network
25. Peter Ton, Silvere Tovignan and Simplicie Davo Vodouhe (Mar 2000),
Endosulfan deaths and poisonings in Benin, *Pesticide News* No. 47. (The Journal of Pesticide Action Network UK)
- 26 Suredranath C, (Aug 15 2001)
"Toxic Tales From God's Own Country", *Down to Earth(English Mag.)*, Vol 10 No.6
- 26a. Sopan Joshi , (Feb 28, 2001)
"Children of Endosulfan", *Down to Earth(English Mag.)*, Vol 9, No.19
27. Thanal Conservation Action and Information Network, Thiruvananthapuram, keralam, India (Oct 2001)
Preliminary findings of the survey on the impact of aerial spraying on the people and the ecosystem: Long Term Monitoring – The Impact of Pesticides on the People and Ecosystem (LMIPPE)
28. Anon (Feb 2001),
Endosulfan Fact sheet (ToxFAQs) Agency for Toxic Substances and Disease Registry (ATSDR), US Dept of Health and Human Services, Public Health Services, Division of Toxicology, Atlanta Georgia
29. T Kannupandi, K Pasupathy, Soundrapandian P, (2000),
Acute toxicity Bioassay of Endosulfan HCH, Copper and Zinc in larvae of Mangrove Crab - *Macrophthalmus erato* (de Man), *J. Mar. Biol. Ass. India*; 42(1&2), 173-177
30. Susan Sang and Sania Petrovic (1999)
Endosulfan- A Review of its Toxicity and its Effects on the Endocrine System WWF (World Wild Life Fund – Canada)
31. Padma S Vankar, Rashmi Mishra, Sapna Johnson (Feb 21, 2001),
Analysis of Samples from Padre Village from Kasargod District of Keralam for Endosulfan Residues (Pollution Monitoring Laboratory- Pesticide Residue Monitoring Study CSE/PRM-1/2001). Centre for Science and Environment, New Delhi.
32. PANUPS
Endosulfan Residue in Australian Beef (*Agro: World Crop Protection News*, Aug 28- 1998, Jan 15 1998); PANUPS (Pesticide Action Network- Update Service- May 20 1996)
33. Anon (Sept 2000)
Toxicological Profile for Endosulfan ATSDR- US Dept of Health and Human Services, Public Health Services.
34. Anon (1975)
Data Sheets on Pesticides No.15- Endosulfan INCHEM- IPCS (International Programme on Chemical Safety)
35. Colborn T, Soto A M, Vomsaal F S (1993)

Developmental Effects of Endocrine Disrupting Chemicals on Wild Life and Humans; Environmental Health Perspectives: 101, 378-384.

36. Hunn J D, Muller E P, De Felice MS (Oct 1993) ' Fish and Agrichemicals- Safeguarding your Pond' GO – 1912; University Extension, University of Missouri – Columbia, Columbia.

37. Anon (1993)
Toxicological profile for Endosulfan – Endosulfan Factsheet – ATSDR, US Dept of Health and Human Services, Public Health Service

38. P Kavitha, A Amruth, K Muthukumaravel
Individual and Synergetic effects of endosulfan and BHC Pesticides on Biochemical Constituents of Aquatic insect (*Sphaerodema rusticum*) Journal of Environment and Pollution 8(2); 201-203.

39. Dalsenti P R, Dallegrove E, Mello J R, Langloh A, Oliveira T R, Faqi A S.(Sept 1999)
Reproductive effects of endosulfan on male offspring of rats exposed during pregnancy and lactation; Hum. Exp. Toxicol., 18(9), 583-589.

40. Tangredi B P, Evans R H (March 1997)
Organochlorine Pesticides Associated with ocular, nasal, or otic infection in the eastern box turtle. (*Terrapene carolina carolina*), J. Zoo. Wildlife. Med.; 28(1), 97-100.

41. Saleh M, Kamal A, Rajab A, El Baro, El Sabae A K (March 1996)
Regional distribution of organochlorine insecticide residues in human milk from Egypt. J. Environ. Sci. Health. B- 31(2); 241-255.

42. Martinez Vidal J L, Moreno Frias M, Garrido Frenich A, Olea Serrano, Olea N (Apr 2000)
Determination of Endocrine-disrupting pesticides and polychlorinated biphenyls in human serum by GC-ECD and GC-MS-MS and evaluation of contribution to the uncertainty of results. Anal. Bioanal. Chem. , 372(7-8), 766- 775.

43. Anderson H R, Vinggaard A M, Rasmussen T H, Gjermansen I M, Bonefeld, Jorgensen E C (Feb 15, 2000)
Effects of currently used pesticides in assays for estrogenicity, androgenicity and aromatase activity invtro. Toxicol. Appl. Pharmacol. , 179 (1); 1-12.

44. Campoy C, Jimenez M, Olea Serrano M F, Olea N, Moreno Frias M, Canabate F, Bayes R, Molina Font J E (Nov 2001)
Analysis of organochlorine pesticides in human milk- preliminary results Early Human Dev. 65 Suppl. S 183- 190.

45. Ntow J (May 2001)

Organochlorine Pesticides in Water, Sediments, crops and human fluids in a farming community in Ghana. Arc. Env. Contam. Toxicol. 40(4) ; 557-563.

46. Cooper S P, Burau K, Smith M A, Symanski E, Sweeny A, Robison T, Colt J S, Laster J, Zalm S H (Nov 2001)
Prenatal Exposure to Pesticides- A feasibility Study Among Migrant and seasonal workers; Am. J. Ind. Med. ; 40(5); 578-585.

47. Willey J B, Krone P H, (Sept 2001)
Effect of endosulfan and nonyl phenol on the primordial germ cell population in prenatal zebrafish embryos. Aquat. Toxicol. 54(1-2); 113-123.

48. Sparling D W, Fellers G M, Mc Connell L L (July 2001)
Pesticides and Amphibian Population Declines in California USA; Environ. Toxicol. Chem. 20 (7); 1591-1595.

49. Preziosi P (1998)
Naturally and Anthropogenic Environmental Estrogens – The Scientific Basis for risk Assessment. Endocrine- disruptors as environmental signalers- An Introduction. Pure and Applied Chemistry; Vol 70, No. 9 ;P 1617-1631.

50. Vengateshwarlu K, Suryarao K, Srinivas V, Sivaprakash K, Jagannadharao N R, Mythilai A (March 2000)
Endosulfan Poisoning- A Clinical Profile, J. Assoc. Physicians India; 48(3); 323-325.

51. Elizen G W (Feb 2001)
Lethal and Sublethal effects of insecticide residues on *Orius insidiosus* (Hemiptera, Anthocoridae) and *Geocoris punctipes* (Hemiptera, Lygaeidae) Journal of Economic Entomology 94(1); 55-59

52. Wirth E F, Lund S A, Fulton M H, Scott G I (June 2001)
Determination of acute mortality in adults and sublethal embryo responses of *Palaemonetes pugio* to endosulfan and methoprene exposure. Aquat. Toxicol. 53 (1); 9-18.

53. Menone M L, Aizpun de Moreno J E, Moreno V J, Lan Franchis AI, Metcalfe T L, Metcalfe C D (Feb 2000)
Arch. Env. Contam. Toxicol. 38(2); 202-208.

54. Park D, Hempleman S C, Propper C R (July 2001)
Endosulfan exposure disrupts Pheromonal system in the red spotted Newt- A Mechanism for subtle effects of environmental chemicals. Environmental Health Perspectives 109(7); 669-673.

55. Younglai E V, Foster W G, Hughes E G, Trimk, Jassel J F (July 2002)
Levels of Environmental Contaminants in Human Follicular Fluid, serum and seminal uple undergoing invitro fertilization: Arch. Environ. Contam. Toxicol., 43(1); 121-126.
56. Anon (2002)
End of the Road for Endosulfan- A Call for Action Against A Dangerous Pesticide- Environmental Justice Foundation, London, UK. Internet site- www.ejfoundation.org
57. Anon (2002)
Death in Small Doses; Combodia's Pesticide problems and Solutions- A report by Environmental Justice Foundation, London, UK . Internet site- www.ejfoundation.org.
58. Chowdhari D K, Nazir A, Saxena D K (2001)
Effect of Three Chlorinated Pesticides on hromosome stress gene in transgenic Drosophila melanogaster. J. Biochem. Mol. Toxicol.15(4); 173-186.
59. Rao A S, Pillala R R (July 2001)
The Concentrations of Pesticides in Sediments from Kolleru Lake in India. Pest Management Science; 57(7), 620-624.
60. Romeo F Quijano (2002)
Endosulfan Poisoning in Kasargod, Kerala, India - A Report on a Fact-Finding Mission, Pesticide Action Network-Asia and the Pacific, Penang, Malaysia.
61. Demeter J, Heyndrickx A, Timperman J, Lefevere M, DeBeer J (July 1977)
Toxicological Analysis in a Case of Endosulfan Suicide Bull. Environ. Contam. Toxicol. 18(1); 110-114.
62. Terry Gips (1990)
Breaking the Pesticide Habit- Alternatives to Twelve Hazardous Pesticides. International Organisation of Consumers Union, Penang, Malaysia. P 348.
63. Anon (2003)
Integrated Management of Coffee Berry Borer(Leaflet), Coffee Board, Division of Entomology/Nematology, Central Coffee Research Insitute, Coffee Research Station, Chikmangalur District, Karnataka, India.
64. Anon(2002)
Final Report of the Investigation of Unusual Illnesses allegedly produced by endosulfan exposure in Padre village of Kasaragod District (N.Kerala); National Institute of Occupational Health, Indian Council for Medical Research, Ahmedabad, India.
65. Vonier P M, D.DA. Crain, J A Mc Lachlan, L J Guillette Jr., Arnold S F (1996)
Interraction of Environmental chemicals with the estrogen and progesteron receptors from the oviduct of the American Alligator. Environmental Health Perstpectives 104; 1318-1322
66. Anon (Nov 12 2001)
World Wildlife Fund Comments on reregistration of Endosulfan submitted to Public Information and Record Integrity Branch, Information Resources and Services Division (7502c) Office of Pesticide Programme, Environmental Protection Agency, Washington.
67. Fransson-Steen R, Flodstram S, Warngard L (1992)
Insecticide Endosulfan and its two stereo isomers promote the growth of altered hepatic foci in rats, Carcinogenesis Vol. 13; ISS 12; P 2299-2303.
- 67a. Clarke, Harvey and Humphrey (1981)
Veterinary Toxicology (II Edition), London.
68. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Arctic Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
69. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Sub Saharan Africa- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
70. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Mediterranean- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
71. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- South East Asia and South Pacific- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
72. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Eastern and Western South America- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
73. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Central America and the Caribbean- Regional Report – Chemicals- United Nations

Environmental Programme- Global Environment Facility

74. Anon (Dec 2002)

Regional Based Assessment of Persistent Toxic Substances- North America- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility

75. Anon (Dec 2002)

Regional Based Assessment of Persistent Toxic Substances- Europe- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility

76. Zou E, Fingerman N (1999)

Effect of Estrogenic Agents on Chitobiase activity in the epidermis and hepato pancrease of the Fiddler Crab (*Uca pugilator*); *EcoToxicol. Environ. Saf.* ; Vol.42;Iss-2; P 185-190.

76a. Zou E, Fingerman N (April 1997)

Synthetic estrogenic Agents do not Interfere with sex determination but do inhibit moulting of the Cladoceran, *Daphnia magna*; *Bull. Environ. Contam. Toxicol.*; 58(4); 594-602.

77. Cerkezayabekir A, Aktae T (1997)

The Histopathological effects of Endosulfan on the Mouse Thyroid Gland; *Turkish Journal of Biology*; 21 ;P 439-444.

78. Anon (2001)

PAN Pesticide Database – Aquatic Ecotoxicity Studies for Endosulfan. (Derived from USEPA AQUIRE Acute Summaries) Pesticide Action Network, Penang, Malaysia.

79. Ciscato C H, Gebara A B, Spinosa Hde S (July 2002)

Pesticides residues in Cow's milk consumed in SaoPaulo City, Brazil; *J. Environ. Sci. Health B.*; 37(4); 323-330.

80. Anon (June 2003)

Pesticide News No 60, The Journal of Pesticide Action Network UK . (Quarterly), P 19.

81. Anon (2003)

Toxicological Profile for Endosulfan ATSDR- US Dept of Health and Human Services, Public Health Services. Web site - <http://www.atsdr.cdc.gov/toxprofiles/tp41-c4.pdf>

82. Mercy Mathew, Andrews M I (Jan 2000)

Effect of Endosulfan on the hatching rate and larval survival of Common Indian Toad (*Bufo melanostictus* Schneider), Proceedings of the 12th Kerala Science Congress, Government of Kerala, pp 615-17

83. Anon (1999)

Global Pesticide Campaigner (Volume 09, Number 2), August 1999. (Source: Latin American News Service, April 4, 1999; [http:// www.lattn.com/cgi-local/daily.pl?094-09.](http://www.lattn.com/cgi-local/daily.pl?094-09.))

84. Devakumar C (Dec 2002)

Endosulfan Aerial Spray Controversy in Kerala. *Pesticide Research Journal*, Vol 14(2): pp:343-344, Society of Pesticide Science, India, Div.of Agricultural Chemicals, IARI, N.Delhi.

85. Anon (Dec 2002)

From the Editors Diary - The High Court Judgement on Endosulfan Episode of Kerala - Excerpts. *Pesticide Research Journal*, Vol 14(2): pp: 350-354, Society of Pesticide Science, India, Div.of Agrichem, IARI, N.Delhi.

86. Ghadiri H and Rose C W (2001)

Degradation of endosulfan in a clay soil from cotton farms of Western Queensland, *Journal of Environmental Management*, 62, pp155-169