

ENDOSULFAN

**Factsheet
and
Answers to Common Questions**

IPEN

Pesticide Working Group

IPEN Pesticide Working Group Project 2004

Endosulfan - Fact sheet and Answers to Common Questions
Usha S. and Harikrishnan V. R.

Design & Lay out: Shibu K. Nair
Printed at Arsha Printers, Thiruvananthapuram
1000 Copies
September 2005

IPEN



International POPs Elimination Network (IPEN) is a global network of public interest non-governmental organizations united to work for the global elimination of Persistent Organic Pollutants (POPs), on an expedited yet socially equitable basis.

Thanal is a public-interest research, advocacy and action organisation based in Kerala, India. It is the International Secretariat for the Pesticide Working Group of IPEN.

IPEN Pesticide Working Group Secretariat,

C/o Thanal, L-14, Jawahar Nagar, Kowdiar, Thiruvananthapuram – 695 003
Kerala, India. Ph : ++91-471-5543001, Fax : ++91-471-2727150
Email : ipenpwg@thanal.org, website : www.thanal.org

Introduction

Endosulfan is a pesticide belonging to the organochlorine group of pesticides, under the Cyclodiene subgroup. Introduced in the 1950's, it emerged as a leading chemical used against a broad spectrum of insects and mites in agriculture and allied sectors. It acts as contact and stomach poison and has a slight fumigant action¹. It is used in vegetables, fruits, paddy, cotton, cashew, tea, coffee, tobacco and timber crops^{1,3,2}. It is also used as a wood preservative and to control tse-tse flies and termites⁵. It is not recommended for household use. Intentional misuse of endosulfan for killing fish^{2,5,7} and snails^{7,1} has also been reported. Endosulfan was also reported as used deliberately as a method of removing unwanted fish from lakes before restoring.^{1,3,2}

Endosulfan was introduced at a time when environmental awareness and knowledge about the environmental fate and toxicology of such chemicals were low and not mandatory as per national laws. But now it is being detected as an important cause of pesticide poisoning in many countries.

Chemical Name

6,7,8,9,10,10- hexa chloro- 1,5,5a,6,9,9a- hexahydro – 6,9- methano- 2,4,3- benzodioxathiepine-3-oxide.


Chemical Formula



In pure form endosulfan exists as colourless crystals. But the technical product is brownish crystals with slight odour of sulphur dioxide³. Technically endosulfan is a mixture of two isomers - alpha-endosulfan and beta-endosulfan in the ratio 7:3. Technical grade endosulfan contains 94% alpha-endosulfan and beta-endosulfan and other related compounds like endosulfan alcohol, endosulfan ether and endosulfan sulfate. Endosulfan is only very slightly soluble in water, but it dissolves readily in xylene, chloroform, kerosene and most organic solvents and is a noncombustible solid. It is mixable with most fungicides and compatible with most pesticides⁴.

Toxicity Classification

The U S Environmental Protection Agency (EPA) classifies endosulfan as Category Ib – Highly Hazardous. The European Union also rates it Highly Hazardous. World Health Organisation (WHO) classifies endosulfan in Category II - Moderately Hazardous. The



classification of WHO was found to be inappropriate considering the classification followed in many countries and the available toxicological information. It has been alleged that the classification is based mainly on LD50 value for acute toxicity generated by the producer company¹³. The Industrial Toxicological Research Centre (ITRC) in India the nodal centre for the Regional Based Assessment of Persistent Toxic Substances (PTS) for the Indian Ocean region by the United Nations Environment Programme-Global Environment Facility(UNEP-GEF) classifies endosulfan as Extremely Hazardous⁴.

Q. What do international treaties say about endosulfan ?

Endosulfan is widely considered to be a Persistent Organic Pollutant (POP) but was not included in the initial list targeted for phase out under the Stockholm Convention. Endosulfan was in the initial list of POPs being considered for world-wide elimination at the first meeting of experts in Vancouver, Canada (1994) jointly convened by governments of Canada and Philippines but was later removed from the list¹³. However, endosulfan is listed as a POP in the Convention on Long-range Transboundary Air Pollution (LRTAP). Endosulfan is recognized as a Persistent Toxic Substance (PTS) by the UNEP.²

Q. What is the regulatory status of endosulfan in world nations ?

Endosulfan has been in world-wide use since its introduction in the 1950's. It was considered a safer alternative to other organochlorine pesticides in many countries in all regions since the 1970's. But in the last two decades many countries have recognized the hazards of wide application of this pesticide and have banned or restricted its use.

Countries which have banned include Singapore, Belize, Tonga, Syria, Germany, Sweden, Philippines, Netherlands, St. Lucia, Columbia, Cambodia, Bahrain, Kuwait, Oman, Qatar, Saudi Arabia, United Arab Emirates, Sri Lanka and Pakistan^{2,73}. Restricted use is allowed in Australia, Bangladesh, Indonesia, Iran, Japan, Korea, Kazakhstan, Lithuania, Thailand, Taiwan, Denmark, Serbia & Montenegro, Norway, Finland, Russia, Venezuela, Dominican Republic, Honduras, Panama, Iceland, Canada, the United States and the United Kingdom ^{2,56,73,80}.

It is one among the twenty one priority compounds identified by the UNEP-GEF in the Regional Based Assessment of Persistent Toxic Substances (PTS),2002. These reports have taken into account the magnitude of use, environmental levels and human and ecological effects of these compounds². In the Indian Ocean Region endosulfan is banned in 8 nations. India is one of the major Indian Ocean rim nations, which has imposed no ban or restrictions on endosulfan. A ban on endosulfan exists in the South Indian state of Kerala (imposed through a Court Order) , which came as a

result of a public pressure following the poisoning of many villages due to aerial spraying of the chemical⁸⁴. Colombia and Cambodia are two countries where endosulfan is banned from use recently⁸⁰.

In the Reports of the Regional Based Assessment of PTS of the UNEP- GEF, endosulfan has been rated depending on the level of concern in the respective region countries. They are as follows-

- ☠ **Indian Ocean region** – “Regional Concern”⁷²
- ☠ **North American region** – Treated as “Regionally specific PTS”, receiving great attention along with HCH, chlordane, PCB, PAHs⁷⁴.
- ☠ **Mediterranean region** – “Local Concern”⁷⁰.
- ☠ **Sub Saharan Africa** – Ranked as the PTS of highest concern after DDT⁶⁹.
- ☠ **Eastern and Western South America** – “Potentially relevant PTS of emerging concern”⁷².
- ☠ **European region** – “Proposed Possible Priority Hazardous Substance”⁷⁵.
- ☠ **South East Asia and Southern Pacific region** – “Regional Concern”. It is identified as a major PTS which has a continuing effect on the natural ecosystem in the region and long term effect on the structure of aquatic eco system⁷¹.
- ☠ **Central America and the Caribbean** – Considered as one of most important PTS of emerging concern⁷³.

Q. How much endosulfan is produced, used and stockpiled ? and Where ?

Practically very little information is available of the volumes of production of endosulfan. WHO estimated that the worldwide production of endosulfan was 10,000 MT in 1984⁸¹. Current estimates of worldwide production or domestic formulations could not be located⁸¹. Many countries that produced endosulfan in the 1970's and 80's do not produce it any more. For instance endosulfan has not been produced in the United States since 1982, but it is still used in chemical formulations⁸¹. In USA, annual average amount of 626 tonnes is used¹³². In European Union, Germany is the only producer of endosulfan and it is manufactured at a single site, where 5000 TPA is produced. Vast majority of this is exported to South America and South East Asia¹³². In EU(1999), the 90% of the endosulfan were used in Mediterranean area¹³². Spain, Italy, Greece and France are the major consumers. Endosulfan is produced mainly in Israel, India, China and South Korea¹³².

India is one of the major producers of endosulfan. Since 1996-97 it produces an average of 8206 MTPA totaling 41033 MT between 1995-2000². India exported

12180 MT during this period and consumed on an average 3599 MTPA. The UNEP-GEF report on PTS has identified some of the producing and importing countries but there is a large data gap. No information regarding stockpiles of endosulfan could be located.

Q. How might I be exposed to endosulfan ?

- § By breathing air near where it has been sprayed.
- § Drinking water contaminated directly or through run-offs.
- § Being in contact with contaminated soil.
- § Eating contaminated food.
- § Smoking cigarettes made from endosulfan contaminated tobacco.
- § Working at endosulfan production centers.
- § Direct ingestion.

Q. How does endosulfan affect human beings ?

Worldwide use of endosulfan increased with the ban/restriction in use of the more persistent organochlorine pesticides like DDT and endrin. Endosulfan is acutely toxic and has been implicated in many cases of poisoning and fatalities. It has been identified with a range of chronic effects, including cancer and impacts on hormonal systems, exhibiting similarities with its predecessors in the organochlorine class.

Acute effects

Endosulfan is highly toxic and can be fatal if inhaled, swallowed or absorbed through the skin. Acute toxicological data is adequately available. Acute oral toxicity is higher than dermal toxicity³⁰. Absorption rate and toxicity is found to increase in the presence of solvents like alcohol and aromatic solvents^{12,24,34,61}.

A number of acute poisoning cases have been reported. Ingestion or breathing high levels of endosulfan may lead to convulsion and death. Endosulfan directly affects the central nervous system^{24,28} and recurrent epileptic seizures are also reported⁹⁴. It is absorbed through skin and eye irritation may also result. Symptoms of poisoning include hyper activity, excitement, dyspnea (breathing difficulty), apnea (stoppage of breathing), salivation, loss of consciousness, diarrhea, anemia, nausea, vomiting, insomnia, blurred vision, cyanosis (bluish discoloration of skin due to want of oxygen), foaming at the mouth, tremor, dry mouth, lack of appetite, irritability, head ache, decreased respiration, loss of memory, haematuria, albuminuria, confusion, dizziness, imbalance and lack of coordination^{4,5,7,30}.

Persons suffering from asthmatic and convulsive disorders are at high risk. Persons on protein deficient diet also possess high risk^{24,30}.

Autopsy examination of an intentional ingestion (suicide) case has revealed damage to liver, lung and brain^{8,8a}.

Chronic effects

There is experimental evidence of adverse effects of endosulfan on the male reproductive system, delaying sexual maturity and interfering with the sex-hormone synthesis¹¹⁸. Endosulfan is a proven endocrine disruptor^{6,9}. It has potential to induce hypothyroidism⁶⁶. Long term health effects are not properly studied, experimented or documented world wide.

Endosulfan exhibits estrogenic properties^{9,43}, comparable to that of DDT⁹. It competes for estradiol for binding to estrogen receptors, thereby inhibiting hormonal function¹⁰⁷. The estrogenic potential of endosulfan increases in the presence of other estrogenic organochlorines¹¹⁰. It induces proliferation of human breast estrogen sensitive MCF7 cells⁴⁹, (*in vitro*) thereby increasing breast cancer risk¹¹⁷. It harms the reproductive system by affecting semen quality, sperm count, spermatogonial cells, sperm morphology and other defects in male sex hormones¹¹. Endosulfan has the capacity to alter the genetic material particularly chromosomes in mammalian cultures¹¹⁹. It is found to inhibit testicular androgen biosynthesis in lab animal experiments³⁰ and exhibits significant risk in renal and testicular damage. It may have adverse effects on central nervous system by inhibiting brain acetyl cholinesterase¹⁶, causing uncontrolled discharge of acetyl choline. Endosulfan ingestion is known to affect the kidneys and liver⁵⁰. It inhibits leucocyte and macrophage migration (this is the inhibition of the natural immune system by disrupting anti-body protection) causing adverse effects on humoral and cell-mediated immune system³⁰. It is also a potential tumor promoter⁶⁷.

Many studies related to its acute and chronic toxicity in laboratory animals are available. Endosulfan is highly toxic to rats and mice^{13, 30}. Some studies suggests its teratogenic²⁸ and carcinogenic properties²¹ on rats and mice. It directly affects the central nervous system, causes liver and kidney (chronic glomerulonephrosis) damage⁶ in rats and mice. It also impairs the reproductive system of rats³⁹. Behavioural and neurological changes have also been noticed³⁰. Thyroid follicular damage in mouse has been reported^{30,66}. Endosulfan is known to damage the endocrine system, nervous system, circulatory, reproductive, respiratory and excretory systems and developing foetus.^{6,7,14,15,16,21,30,37}

The National Institute of Occupational Health (India) have linked the higher prevalence of neurobehavioral disorders, congenital malformations in female children and abnormalities related to male reproductive systems to the continuous exposure to endosulfan spray. The study was conducted among children in one of the villages in Kasaragod District (in the South Indian State of Kerala) where endosulfan was aerially sprayed⁶⁴. Endosulfan is

implicated in the occurrence of adverse health effects particularly in rural communities in South East Asia, Southern Pacific and Sub-Saharan Africa.^{69,71}

Q. Can endosulfan cause cancer ?

Endosulfan is found to damage human red blood cells (RBC) at concentrations of 1ppb-1ppm¹⁰. Both alpha-endosulfan and beta- endosulfan are genotoxic to HepG2 cells⁸. Endosulfan is hepatotoxic⁶, mutagenic, clastogenic and induce effects on cell cycle kinetics¹³. Endosulfan has been shown to cause chromosomal aberrations in hamster and mouse and sex linked mutations in *Drosophila*^{13,30,58}. Endosulfan has caused mutations in bacterial and yeast cell. It is also known to cause mutations in mammals¹³

A re-analysis data from a 1978 NCI (National Cancer Institute, US) study in Osborne-Mendel rats has revealed that endosulfan induced malignant neoplasms at all sites in male and female rats and endocrine organs in male rats²¹. Both sexes developed lymphosarcomas and female rats had neoplasms of the reproductive system²¹. Endosulfan is also carcinogenic for the liver of female mice²¹.

No accurate data related to the carcinogenicity of endosulfan in human is available but from field level reports, endosulfan can be highly suspected for having carcinogenic properties in human beings, especially in cases of chronic exposure¹³. In some reports it is referred to as having possible carcinogenic effects, effects in human immune and reproductive system⁶⁸. Studies have also shown that it induces proliferation of human breast estrogen sensitive MCF7 cells in vitro^{9,49} which may lead to greater breast cancer risk. Studies also indicate the contribution of endosulfan in the combined effect of environmental estrogens in inducing breast cancer¹¹⁷.

Q. What happens to endosulfan in the environment ?

The fate of endosulfan released in the environment is different for the two isomers and also depends on the medium it gets deposited.

Beta- endosulfan is more persistent than its alpha- isomer. Endosulfan sulphate is the main degradation product of both isomers, which is equally toxic^{27,66} and is itself more persistent in the environment than its parent compounds^{5,30}. Endosulfan can be broken down by photolysis, hydrolysis and bio degradation. Endosulfan diol, endosulfan lactone, endosulfan ether etc. are some of the other byproducts⁵. Although the isomers are fairly resistant to photolysis, the break down products are susceptible¹³². On plant surfaces endosulfan rapidly degrades to metabolites³⁰. Endosulfan is fairly immobile in soil and is highly persistent^{13,33}. Major products in soil are endosulfan diol, endosulfan sulfate and endosulfan lactone. Endosulfan sulfate production increases with an increase in

temperature³⁰. Endosulfan will persist longer under more acidic conditions^{5,24}. It persists longer under submerged conditions⁸⁶. The half life of endosulfan varies from 60 days (alpha- endosulfan) to 800 days (beta-endosulfan)¹³. It enters air by volatilization from plant and water surfaces^{13,33}. Contamination by drift and particle transport also occur³³. Ultra low volume (ULV)- application can drift several kilometers from point of application³⁰. It has been detected in remote areas including the Arctic¹³² in air, snow-water and lake waters, rainfall and snow samples in Californian mountains and remote European mountain lakes^{5,30,33,68,75,132}. Beta-endosulfan is more stable in air. Endosulfan has been shown to be released from wood preservatives into room atmosphere over an one year period of observation⁵. In water endosulfan has a half life of 35 to 150 days¹³. It does not easily dissolve in water and may accumulate in bodies of fish and other aquatic organisms^{30,69}. The break down products in water are endosulfan-diol and endosulfan sulfate. Endosulfan does not reach down much to ground water³⁰, but has been proven to run off after spraying. But it has been detected from ground water at deep soil layers in concentration ranging from 0.009- 0.053 micro gram per litre up to 20 days after last spraying¹³. USEPA recommends that the levels of endosulfan in rivers, lakes and streams should not be more than 74 ppb²⁸. But this limit is 15 times more than the concentration required for causing reproductive damage in red spotted newt⁵⁴. Several studies of bioaccumulation have been conducted around the world and residues were found in aquatic organisms. USEPA considers endosulfan as having a high bioaccumulation potential in fish¹⁶, but not much evidence of bioaccumulation in higher trophic levels is available.

Q. Is there any evidence of endosulfan contamination in the environment, food and human beings ?

Reports of endosulfan residue in food, soil, air, body tissues etc are available from all parts of the globe.

Residues were detected from air, water (surface and ground water)^{2,59} and soil in India², water and sediments in Ghana⁴⁵, marine water and sediments in India^{2,87}, shallow ground water in Pakistan¹²⁴, river water in China¹²⁷, lagoons in Spain, surface and ground waters in Portugal⁹², ground and well waters in the Philippines^{133,134}, coastal, estuarine and river sediments in Israel (high concentrations in Lake Kinneret)⁷⁰, water in Benin, Malawi, Nigeria⁶⁹ and from drinking, ground, surface and marine waters in South Africa⁹⁷ (alarming levels in river water 684-4843 ng/L)⁶⁹, soil in Benin, Nigeria, Sudan and Zambia, sediments in Benin and Nigeria, vegetation in Madagascar, Zambia and Ghana^{69,97}, Paddy fields in Mediterranean¹³¹, water from remote mountain lakes in Europe (the Alps, Caledonian and the Pyrenees)⁷⁵ and river and sea water in South East Asia⁷¹ etc. were found to be polluted with endosulfan. In Malaysia river sediments were found to have high levels (434micro grams/litre) of endosulfan⁷¹. Reports from Central America shows

air and wetlands⁹³ in Costa Rica, shallow lakes, coastal waters, estuaries, well, surface and marine waters and sediments in Honduras, Mexico, Argentina and Jamaica are contaminated with endosulfan residues^{73,96,113,130}. Endosulfan is one of the most frequently reported PTS in surface and ground water in Central America and the Caribbean⁷³. Toxic levels of endosulfan are reported from the coastal lagoons of Mexico¹⁰⁵. In Guatemala, it is found more frequently and at higher levels in river water, well water and spring water⁷³. It has been detected in municipal water system in Colombia⁷³. Endosulfan residues were found in high levels from the gulf of Mexico and from many parts of the USA¹⁰⁴. Residues were also found in the Greenland biota¹¹⁵ and biota including mammals of Arctic region¹³².

High concentrations of alpha and beta endosulfan isomers and endosulfan sulphate have been detected in tree bark samples through out the world, particularly in India and the Pacific Rim³⁰.

Endosulfan has been detected from food samples²⁰ around the world- in Australian beef at 0.36 mg/kg³² (2 times the Australian limit and 4 times the international limit), in cows milk from tobacco farming areas in USA and food samples in USA and Canada⁵. Residues were detected in tomatoes from Brazil²⁰, untreated leek in Argentina⁹⁵, Spanish pepper samples from Finland, fish from India², Kenya and Nigeria⁶⁹, food and vegetables in Croatia⁷⁰ and vegetables in Cyprus⁷⁰. Residues were also found from sunflower seeds of untreated areas in Spain¹¹². High levels of residues were found in red pepper and egg plants in Catania(Italy)⁷⁰. High residues have also been found in diary food, meat, fish, chicken, and vegetables in Eastern and Western South America⁷², cows milk in Brazil⁷⁹, vegetable diet in Kuwait, vegetables, vegetable oil and seeds from India and animal samples from slaughter houses in India². Endosulfan residues were detected from cows milk in Colombia far exceeding the WHO and FAO reference levels⁷³. Fish from Kenya and Nigeria were heavily contaminated with endosulfan⁶⁹. The European Union has banned import of fish from Tanzania, Uganda and Kenya due to high levels of endosulfan residues⁸⁰. Residues were also detected from animals samples in Benin, Nigeria, Cote d' Ivoire, Madagascar, South Africa and Kenya⁶⁹. Presence of endosulfan in Lake Trout in North America suggests a wide dispersal from areas of use to isolated lakes⁷⁴.

Endosulfan has also been detected from human tissues. It has been detected from cord blood samples obtained at the time of delivery⁴⁶, human sera^{42,55}, adipose tissue⁸⁷ and human milk samples obtained from healthy lactating women in Spain⁴⁴. Human breast milk from Egypt^{30,41}, Colombia and Nicaragua⁷³ and from cotton pickers in Pakistan² were found to be contaminated. High levels of endosulfan were detected in human breast milk in Sub Saharan Africa⁶⁹ and also from India¹²⁸. Residues were also detected in fat samples from children living nearby farms in Spain²². Blood, human milk and urine samples in Croatia were also found to be contaminated. Alarming high levels of

endosulfan residues were observed in human blood and milk in a study in Kasaragod in Kerala, India^{26,26a,27,31,64}.

Q. How does endosulfan affect wildlife and domestic animals?

There are many reports regarding the toxicity of endosulfan on wildlife. National Wildlife Federation (USA) states that endosulfan is extremely toxic to wildlife and acutely toxic to bees¹³. It is acutely toxic to birds - mallard ducks²⁴, quails and pheasants³⁰. The alpha isomer is more neuro-toxic and its acute toxicity against mammals is more than three times that of the beta-isomer¹⁰⁹. It is genotoxic and is a known endocrine disruptor in terrestrial and aquatic species¹³²

Endosulfan is highly toxic to aquatic organisms even at recommended levels of application^{13,71,88,132}. It is particularly toxic to fishes^{5,18,24,132} – massive fish kills are reported from many places¹³. It also causes endocrine problems, reduction of protein in tissues and other health effects. The high toxicity to fish species is evidenced through studies on *Gambusia affinis*¹⁸, Rainbow trout, channel catfish, bluegill sunfish, minnow^{24,30,36}. It affects metabolism in freshwater fishes by inhibiting transcription at some points¹²⁶. It is known to impair the pheromonal systems leading to disrupted male choice and lowered mating success in female red spotted newts⁵⁴. It is known to affect the germ cell population of zebrafish embryos⁴⁷. It exhibits anti-estrogenic effects in fresh water catfish, reproductive problems in female teleostfish and opercooidfish³⁰. It also bioconcentrates in aquatic organisms³³. Reports from South East Asia and Southern Pacific proves that endosulfan has detrimental effects on aquatic biota⁷¹. Reports prove its high toxicity to frogs, toads, annelids, snails, aquatic insects (damsel flies, midges, beetles etc), crustaceans (crab, shrimps, prawns etc), fishes and molluscs^{78, 114, 125}. Endosulfan affects the hatching rate and larval survival of common Indian Toad⁸². It also affects the larval survival, growth rate, respiration and caused limb deformities in the streamside salamander¹⁰³. Reports from Argentina and South Africa suggests that it affects the aquatic insect and macro-invertebrate populations in streams and rivers^{122,123}. Studies show that it has contributed to the adverse effects on the wetland ecosystem in the Republic of Azherbaijan⁹⁹.

Besides being toxic to crab larvae²⁹, it has been reported to disrupt the moulting of crabs and aquatic invertebrates^{76,76a}. It is known to function as an endocrine disruptor in American alligators⁶⁵. It has been detected from the nonviable eggs of Morelet's crocodile¹⁹. It is also known to be affecting the egg laying of grass shrimps⁵². It has been detected as one of the predominant organochlorine pesticides in the silverside fish in Argentina⁵³. It has been related to the drastic population decline of anuran amphibians in Western USA in over the last 10 to 15 years⁴⁸. Residues have been obtained from the liver of Eastern Box turtle⁴⁰. It is found to accumulate 600 times the water concentration in mussels²⁴.

Endosulfan was found toxic to earthworms, causing a significant reduction in the growth rate and total protein content¹²⁹. It is toxic to non target organisms^{16,120}, like predators of several pests⁵¹. Endosulfan is highly toxic to soil micro arthropods¹⁷,

micro organisms, zooplanktons⁹⁰, phytoplanktons, soil algae, actinomycetes, bacterial colonies etc.^{13,30}. It has shown to cause chromosomal aberrations in *Drosophila* fly¹³². It is also toxic to mammals like rabbits^{13,30} and rats^{21,108}. The disappearance of cats, frogs, bees, fresh water fishes etc. were reported from Kasaragod District in South India, where endosulfan was aerielly sprayed²⁷.

Endosulfan is also highly poisonous to cats (LD50-2mg/kg) and dogs (LD50-76.7mg/kg). Reversible blindness and lack of muscle coordination has been reported in sheep and cattle grazing on endosulfan contaminated grass²⁴. Farmers from endosulfan sprayed area report of acute effects to cattle at time of spray and after, and leading to fatalities²⁷. Acute poisoning of cattle and cattle deaths, due to the direct use of endosulfan on cattle as an ecto-parasitic-control, are reported from Turkey¹⁰⁰. Similar poisoning case is reported from USA also¹⁰¹. Chronic exposure to endosulfan leads to deleterious effects on metabolism and immune system of broiler chicken¹¹¹.

Endosulfan exhibits phytotoxic properties⁷. Toxic effects on plants like root growth inhibition, stunting, burning of tips and margins of leaves and affected root permeability have been reported¹³. It is a prominent contaminant in vascular plants and lichen even in remote areas like the Arctic^{68,115}. It is toxic to fresh water green algae⁹⁰ and also to blue-green algae⁹¹. It also affects diatom abundance, chrysophytes, cryptophytes and dinoflagellates¹²¹.

Q. Are there reports of endosulfan poisoning from around the world ?

Cases of endosulfan poisoning have been reported from many parts of the world. Accidental and intentional exposure leading to human fatalities and environmental tragedies has occurred. The following are some of the major cases of poisoning.

Human poisonings

In Sudan in 1988, endosulfan barrels washed in irrigation canals caused fish kills and three people died after drinking water from the canal¹³. In the Philippines, endosulfan accounted for the largest number of deaths due to pesticide poisoning reported in 1991¹³. In Sulawesi, Indonesia, 32 cases of poisoning due to endosulfan have been reported from 1990 to 1993⁵⁶. In Columbia, 155 cases of poisoning (in 1994) and 60 cases (1993) of poisoning due to endosulfan were reported⁵⁶. In Northern India, 18 cases of endosulfan poisoning have been reported in 1995-97 by accidental over-exposure during spray²³. Endosulfan caused a rise in death numbers due to poisoning in Srilanka from 1994 to 1998⁹⁸. The misuse of endosulfan to kill snails has resulted in the largest number of poisoning cases with fatalities in Philippines in 1996⁷⁵. Poisoning due to consumption of endosulfan-contaminated food is also reported from Turkey (2003)¹⁰². Many cases of poisoning-death in Guatemala, Costa Rica and other Central American

countries have also been reported⁷³. In Feb 2002, two South African boys living near Ntabamhlophe, Kwa-Zulu Natal died following exposure to endosulfan. A police officer and 3 journalists were hospitalized after visiting the place several hours later⁸⁰. In 2000, a case of 44 individuals who consumed food accidentally contaminated by endosulfan was reported in rural India. One individual died in the incident⁴⁴. In 2004, 36 persons of all age groups in a rural area of Jabalpur, India were poisoned after consuming wheat-grains or flour contaminated with endosulfan¹⁰⁶.

***The worst of all the cases so far reported are from three nations—
Cuba, Benin and India.***

Pesticide Poisoning in Cuba : Endosulfan was responsible for the death of 15 people in the western province of Matanzas, Cuba in February 1999. A total of 63 people became ill after consuming food contaminated with endosulfan according to Cuban authorities⁸³.

The Benin Tragedy:- In Borgou province in Benin, endosulfan poisoning caused many deaths during 1999 - 2000 cotton season. Official records state that at least 37 deaths occurred and 36 were taken seriously ill. In the same region in 1999 a boy died after eating corn contaminated with endosulfan²⁵.

The Kasaragod Tragedy, South India:- People in 15 villages in Kasaragod in the South Indian State of Kerala were subjected to continuous exposure to endosulfan which was aerielly sprayed three times every year for 24 years. Congenital Birth defects, reproductive health problems, Cancers, loss of immunity, neurological and mental diseases were reported among the villagers. Following a public outcry a number of health based scientific studies confirmed that the health problems were directly linked to the exposure to endosulfan^{27,60,64}.

Poisoning of water bodies and fish kills

A wide spread fish kill was observed in 1969, when 30 kilograms of endosulfan was discharged into the Rhine river in Federal Republic of Germany^{5,30}. In 1975, an accidental spill of endosulfan caused a major fish kill in North Brook a tributary of the Dunk River in Eastern Prince County of Prince Edward Island. The Brook trout population was reduced from 2227-4147 to 45-246³⁰. In 1995, run-off from cotton fields contaminated with endosulfan resulted in the death of more than 24,000 fish along a 25-kilometer stretch of a river in Alabama¹³. Investigations showed that the pesticide had been sprayed according to label instructions.

Q. What are the alternatives to endosulfan?

While there are many non-chemical alternative practices, the choice of one primarily depends on the crop and the pest for which it has been used.

One of the important crops on which endosulfan is used is cotton. Endosulfan is also used on crops like vegetables, cashew, coffee, tea etc.

There are numerous documented benefits from the adoption of Integrated Pest Management (IPM) programme for cotton from different cotton growing areas all over the world. The focus of IPM is on cultural and biological control of pests rather than depending on pesticides. A comparison in Frio county, Texas show a savings of between 131- 300 percent for alternative cotton pest management⁶². Organic farming of cotton is also gaining popularity among cotton farmers in the Asian region⁶². Similarly in vegetable cultivation many field trials have been carried out on alternative pest control methods focusing on herbal pesticides. In some studies, botanical pesticides were found more effective than endosulfan in controlling green house pests¹¹⁶. In the Asian region farmers have developed their own combinations and methods of pest control using chillies, garlic, asafoetida, cow urine and many other plant materials. FAO supported IPM programme in vegetables is well documented.

In India endosulfan was recommended for controlling tea mosquito bug in cashew and tea until 2000. Following the Kasaragod Tragedy, the National Research Centre for Cashew has withdrawn its recommendation for endosulfan use in cashew. Many cashew farmers have tried organic method and application of neem oil and pongamia leaf extract is found successful by some South Indian farmers. Some farmers have even tried 'do nothing' farming and the results are encouraging. Alternative pesticides and organic farming are encouraged in tea plantations and the use of endosulfan is being eliminated. There are tea plantations in South India which has adopted bio-dynamic farming and have completely eliminated the use of chemical pesticides including endosulfan.

In coffee, to control coffee berry borer, Integrated Pest Management is adopted instead of endosulfan⁶³. This includes strict phyto-sanitary and cultural measures supplemented with bio control measures and mass trapping of the insects.

Though alternatives to endosulfan are available, support for such practices are very low. If Governments and research institutions can support such work, use of endosulfan can be totally eliminated in agriculture and other sectors.

Q. What can I do to eliminate the use of endosulfan?

- ◆ Look out for the alternative non-chemical pest control, which is available in your region and other parts of the world, and thus eliminate the use of endosulfan.

- ◆ Pressure your government to ratify the Stockholm Convention if it has not already done so (see <http://www.pops.int> for a current list) and press for addition of endosulfan to the Stockholm Convention List.
- ◆ Pressure your Government to ban the production and/or use of endosulfan, if it has not already done so.
- ◆ Join one of the local and international campaigns to ban endosulfan.
- ◆ Adopt "Land and Food without Poisons" as a goal for survival and adopt organic, ecological or natural agriculture in your farm. Share this message among fellow farmers.

Trade names and formulations of endosulfan:-

Endosulfan is available in the market in a number of trade names. It is available as formulations of Emulsifiable Concentrate (EC), Wettable Powder (WP), Ultra low Volume Liquid (ULV), Granules (G), Dust(D) and Smoke Tablets.

The following list contains some of its trade names. The list is not complete and some names may now be obsolete.

Bangladesh – Thiodan

Brunei – Thiodan, Fezdion

Chile – Parmazol E, Flavylon, Galgofan, Thiodan, Thionex, Thionyl y methofan

India – Agrosulphan, Agiro Sulphan, Banej Sulphan, Cilo Sulphan, Endo Sulphan, E-Sulfan, Endo Chithin, Endocid, Endonit, Endomil, Endosol, Endostar, Endosun, Endotaf, Endostan, Endocing, Endocide, Endosulpher, Gaydan, Gilnore Endorifan, Hexa-sulfan, Hildan, Hockey Endosulfan, Hy-sulfan, Kemu Sulfan, Hilexute-Sulfan, Krushi Endosulfan, Lusu Sulfan, Marvel-Micosulfan, Mico Thansulfan, Pary Sulfan, Pesticel, Remisfan, Sico sulfan, Solesulfan, Sujadin, Sulfan, Tej Sulfan, Thiodon, Thiokill, Thionel, Thionex, Thioton, Veg-fru Thiotox, Veg-fru Thiotex, Vika sulfan.

Indonesia – Thiodan, Fanodan, Dekasulfan.

Korea – Malix, Thioliex.

Pakistan – Siagon, Thiodan, Thioluxan.

Philippines – Atlas Endosulfan, Endosulfan, Contra, Endox, Thiodan.

Sri Lanka – Thiodan, Thionex, Endomack, Endocel, Bauris Endosulfan, Harcros Harcosan, Red Star Anglo-sulfan.

Thailand – Thiodan, A. B. Fan, Aggrodan, Agridan, Bensodan, Bensocarb, Beosit, Brook, Clement, Dew Dan, Dior 35, Dori, Dumpersan, E C Sulfan, Egodan, Endan, Endodan, Endosulfan, Endrew, Endye, Endyne, Etonic, Exxo-Z, Famcodan, Fortune, Freedan, Gardner, Gycin, Hor Mush, Hydrodan, J-teedan, Jack Dum, Kasidan, L P dan, Lordjim, Malix, Manyoo, Metrodan, Nayam, Newcodan, Nockdyne, Ox Xa, Patodan, Pestdye, Pro-d-dan, Sandan, Shevanex, Simadan, Sonydan, Summer, Tanadan, Teophos, Thanyacarb, Thimul, Thiofor, Torpidan, Urofen, Wephos, Zumic.

Other Names: Chlorthiepin, Cyclodan, Endox, Thifor, Thiomul, Thionate.

References

1. K.K. Nayar, T.N Ananta Krishnan and David B.V.
General and Applied Entomology. Tata McGraw Hill Publishing Co Ltd, New Delhi, India.
 2. Anon (Dec 2002).
Regional Based Assessment of Persistent Toxic Substances – Indian Ocean Regional Report – Chemicals – United Nations Environmental Programme – Global Environment Facility (UNEP – GEF).
 3. Anon (June 1997).
NIOSH Pocket Guide to Chemical Hazards. US Dept of Health and Human Services – National Institute for Occupational Safety and Health.
 4. Anon (1989).
Toxicity Data Hand Book. Vol:III, Pesticide-A. Industrial Toxicology Research Centre (Council of Scientific and Industrial Research) Lucknow, India.
 5. Anon (1984).
Environment Health Criteria 40- Endosulfan. IPCS (International Programme on Chemical Safety) – WHO Geneva.
 6. Anon (22nd Dec 1997).
National Registration Authority for Agriculture and Veterinary Chemicals, Australia. Existing Chemical Review Programme – NRA Review of endosulfan – Draft.
 7. Anon (Issue 2 May, 2000)
Hazardous Substances Data Book (HSDB)- US National Library of Media – Canadian Centre for Occupational Health and Safety.
 8. Yuquan Lu, Kane Hiza MoriMoto, Tatsuya Takeshita, Toru Takeuchi, Takeshi Saito.
Genotoxic effects of A-Endosulfan & B-Endosulfan on Human HEPG2 cells – Environmental Health Perspectives ; JUNE 2000 ; 108-(6), 559-561
 - 8a. Boerehoom F T, Van Dijk A, Van Zoonan P, Meulenbelt J (1998)
Nonaccidental Endosulfan Intoxication, A case report with toxico kinetic Calculations and Tissue concentration. J Toxicol. Clin Toxicol , 36: 345-352.
 9. Soto AM; Chung K L, Sonnen Schein C(1994).
The Pesticides Endosulfan, Toxaphene and Dieldrin have estrogenic effect on human estrogen – sensitive cells. Environmental Health Perspectives – Vol 102, Iss 4; P 380-3.
 10. Daniel C S, Agarwal S, Agarwal SS(1986)
Human Red blood cell membrane damage by Endosulfan, Toxicol. Lett; Vol.32, ISS !-2, P 113-8. General and Applied Entomology. Tata McGraw Hill Publishing Co Ltd, New Delhi, India
 11. Pandey N, Gundevia F, Prem A S, Ray P K. (1990)
Studies on the Genotoxicity of Endosulfan, an organochlorine insecticide in mammalian germ cells. Mutant Res; Vol. 242; ISS 1, P 1-7.
 12. Fransis A Gunth (1968) (Edit)
Residue Reviews, Vol. 22.
-

13. Romeo F. Quijano, MD (Oct/Dec 2000).
Risk Assessment in a third world reality: An Endosulfan case History. International Journal of Occupational and Environment Health. Vol. 6, No. 4.
14. Paul V, Balasubrahmaniam E, Jayakumar A R, Kazi M. (1995)
A sex related difference in the neuro behavioural and hepatic effects following chronic endosulfan treatment in rats. Eur. J. Pharmacol; Vol.293, ISS 4, P 355-60
15. Sinha N, Narayan R, Sankar R, Saxena D K. (1995).
Endosulfan induced bio chemical changes in testis of rats. Vet. Human Toxicol; Vol. 37, ISS 6, P 547-9.
16. Naqvi. S M, Vaishnavi C (1993)
Bio Accumulative Potential and toxicity of Endosulfan insecticide to non target animals. Comp. Biochem. Physiol. C; Vol 105, Iss 3, P 347-61.
17. Joy V C, Chakravarthy P P (1991)
Impact of Insecticide on Non Target Micro Arthropod Fauna in Agricultural Soil. Ecotoxicol. Environ. Saf. ; Vol. 22, ISS 1, P 8-16.
18. Karim A. A, Haridi A. A, el Rayah E A (1985)
The environmental impacts of four insecticides on non target organisms in the Gezira Irrigation Scheme canals of Sudan. J. Trop. Medic. Hyg: Vol 88, ISS 2, P 161-8.
19. Wu T. H, Rainwater T. R, Platt S. G, Mc Murry S. T, Anderson T. A (March 2000)
Organo chlorine contaminants in Morelet's crocodile (Crocodylus moreletti) eggs from Belize. Chemosphere; 40(6); 671-8
20. Araujo A C, Telks D L, Gorni R, Lima L L (June 1999)
Endosulfan residues in Brazilian Tomatos and their impact on public health and the environment. Bulletin Environ. Contam. Toxicol; 62(6): 671-6.
21. Reuber MD , (Aug 1981)
The role of toxicity in the carcinogenicity of Endosulfan . Sci. Total Environ.; 20(1); 23-47.
22. Olea- Serrano F, Olea N, Lardelli-Claret P, Rivas A, Barba-Navarro A. Jan-Mar (1999).
Inadvertant exposure to xenoestrogens in children. Toxicol Ind. Health, 15(1-2); 151-8.
23. Chugh.S.N, Dhawan R, Agarwal N, Mohajan S K, (Sept 1998).
Endosulfan poisoning in north India, a report of 18 cases. International Journal Clin. Pharmacol Ther; 36(9); 474-7
24. Anon(1996)
Pesticide information Profile- Endosulfan (Revised June 1996)EXTONET- Extention Toxicology Network
25. Peter Ton, Silvere Tovignan and Simplicie Davo Vodouhe (Mar 2000),
Endosulfan deaths and poisonings in Benin, Pesticide News No. 47. (The Journal of Pesticide Action Network UK)
- 26 Surendranath C, (Aug 15 2001)
"Toxic Tales From God's Own Country", Down to Earth(English Mag.), Vol 10 No.6
- 26a. Sopan Joshi , (Feb 28, 2001)
"Children of Endosulfan", Down to Earth(English Mag.), Vol 9, No.19

27. *Thanal Conservation Action and Information Network, Thiruvananthapuram, Kerala, India (Oct 2001)*
Preliminary findings of the survey on the impact of aerial spraying on the people and the ecosystem: Long Term Monitoring – The Impact of Pesticides on the People and Ecosystem (LMIPPE)
28. *Anon (Feb 2001),*
Endosulfan Fact sheet (ToxFAQs) Agency for Toxic Substances and Disease Registry (ATSDR), US Dept of Health and Human Services, Public Health Services, Division of Toxicology, Atlanta Georgia
29. *T Kannupandi, K Pasupathy, Soundrapandian P, (2000),*
Acute toxicity Bioassay of Endosulfan HCH, Copper and Zinc in larvae of Mangrove Crab -Macrophthalmus erato (de Man), J. Mar. Biol. Ass. India; 42(1&2), 173-177
30. *Susan Sang and Sania Petrovic (1999)*
Endosulfan- A Review of its Toxicity and its Effects on the Endocrine System WWF (World Wild Life Fund – Canada)
31. *Padma S Vankar, Rashmi Mishra, Sapna Johnson (Feb 21, 2001),*
Analysis of Samples from Padre Village from Kasargod District of Kerala for Endosulfan Residues (Pollution Monitoring Laboratory- Pesticide Residue Monitoring Study CSE/PRM-1/2001). Centre for Science and Environment, New Delhi.
32. *PANUPS*
Endosulfan Residue in Australian Beef (Agro: World Crop Protection News,Aug 28-1998, Jan 15 1998); PANUPS (Pesticide Action Network- Update Service- May 20 1996)
33. *Anon (Sept 2000)*
Toxicological Profile for Endosulfan ATSDR- US Dept of Health and Human Services, Public Health Services.
34. *Anon (1975)*
Data Sheets on Pesticides No.15- Endosulfan INCHEM- IPCS (International Programme on Chemical Safety)
35. *Colborn T, Soto A M, Vomsaal F S (1993)*
Developmental Effects of Endocrine Disrupting Chemicals on Wild Life and Humans; Environmental Health Perspectives: 101, 378-384.
36. *Hunn J D, Muller E P, De Felice MS (Oct 1993)*
Fish and Agrichemicals- Safeguarding your Pond, GO – 1912; University Extension, University of Missouri – Columbia, Columbia.
37. *Anon (1993)*
Toxicological profile for Endosulfan – Endosulfan Factsheet – ATSDR, US Dept of Health and Human Services, Public Health Service
38. *P Kavitha, A Amruth, K Muthukumaravel*
Individual and Synergetic effects of endosulfan and BHC Pesticides on Biochemical Constituents of Aquatic insect (Sphaerodema rusticum) Journal of Environment and Pollution 8(2); 201-203.
39. *Dalsenti P R, Dallegrave E, Mello J R, Langloh A, Oliveira T R, Faqi A S.(Sept 1999)*
Reproductive effects of endosulfan on male offspring of rats exposed during pregnancy and lactation; Hum. Exp. Toxicol., 18(9), 583-589.
-

40. Tangredi B P, Evans R H (March 1997)
Organochlorine Pesticides Associated with ocular, nasal, or otic infection in the Eastern box turtle. (*Terrapene carolina carolina*), *J. Zoo. Wildlife. Med.*; 28(1), 97-100.
41. Saleh M, Kamal A, Rajab A, El Baro, El Sabae A K (March 1996)
Regional distribution of organochlorine insecticide residues in human milk from Egypt. *J. Environ. Sci. Health. B-* 31(2); 241-255.
42. Martinez Vidal J L, Moreno Frias M, Garrido Frenich A, Olea Serrano, Olea N (Apr 2000)
Determination of Endocrine-disrupting pesticides and polychlorinated biphenyls in human serum by GC-ECD and GC-MS-MS and evaluation of contribution to the uncertainty of results. *Anal. Bioanal. Chem.* , 372(7-8), 766- 775.
43. Anderson H R, Vinggaard A M, Rasmussen T H, Gjermansen I M, Bonefeld, Jorgensen E C (Feb 15, 2000)
Effects of currently used pesticides in assays for estrogenicity, androgenicity and aromatase activity invitro. *Toxicol. Appl. Pharmacol.* , 179 (1); 1-12.
44. Campoy C, Jimenez M, Olea Serrano M F, Olea N, Moreno Frias M, Canabate F, Bayes R, Molina Font J E (Nov 2001)
Analysis of organochlorine pesticides in human milk- preliminary results *Early Human Dev.* 65 Suppl. S 183- 190.
45. Ntow J (May 2001)
Organochlorine Pesticides in Water, Sediments, crops and human fluids in a farming community in Ghana. *Arc. Env. Contam. Toxicol.* 40(4) ; 557-563.
46. Cooper S P, Burau K, Smith M A, Symanski E, Sweeny A, Robison T, Colt J S, Laster J, Zalm S H (Nov 2001)
Prenatal Exposure to Pesticides- A feasibility Study Among Migrant and seasonal workers; *Am. J. Ind. Med.* ; 40(5); 578-585.
47. Willey J B, Krone P H, (Sept 2001)
Effect of endosulfan and nonyl phenol on the primordial germ cell population in prenatal zebrafish embryos. *Aquat. Toxicol.* 54(1-2); 113-123.
48. Sparling D W, Fellers G M, Mc Connell L L (July 2001)
Pesticides and Amphibian Population Declines in California USA; *Environ. Toxicol. Chem.* 20 (7); 1591-1595.
49. Preziosi P (1998)
Naturally and Anthropogenic Environmental Estrogens – The Scientific Basis for risk Assessment. *Endocrine- disruptors as environmental signalers- An Introduction.* *Pure and Applied Chemistry*; Vol 70, No. 9 ;P 1617-1631.
50. Vengateshwarlu K, Suryarao K, Srinivas V, Sivaprakash K, Jagannadharao N R, Mythilai A (March 2000)
Endosulfan Poisoning- A Clinical Profile, *J. Assoc. Physicians India*; 48(3); 323-325.
51. Elizen G W (Feb 2001)
Lethal and Sublethal effects of insecticide residues on *Orius insidiosus* (Hemiptera, Anthocoridae) and *Geocoris punctipes* (Hemiptera, Lygaeidae) *Journal of Economic Entomology* 94(1); 55-59

52. Wirth E F, Lund S A, Fulton M H, Scott G I (June 2001)
Determination of acute mortality in adults and sublethal embryo responses of *Paleomonetes pugio* to endosulfan and methoprene exposure. *Aquat. Toxicol.* 53 (1); 9-18.
53. Menone M L, Aizpun de Moreno J E, Moreno V J, Lan Franchis Al, Metcalfe T L, Metcalfe C D (Feb 2000)
Arch. Env. Contam. Toxicol. 38(2); 202-208.
54. Park D, Hempleman S C, Propper C R (July 2001)
Endosulfan exposure disrupts Pheromonal system in the red spotted Newt- A Mechanism for subtle effects of environmental chemicals. *Environmental Health Perspectives* 109(7); 669-673.
55. Younglai E V, Foster W G, Hughes E G, Trimk, Jassel J F (July 2002)
Levels of Environmental Contaminants in Human Follicular Fluid, serum and seminal uple undergoing invitro fertilization: *Arch. Environ. Contam. Toxicol.*, 43(1); 121-126.
56. Anon (2002)
End of the Road for Endosulfan- A Call for Action Against A Dangerous Pesticide- Environmental Justice Foundation, London, UK. Internet site- www.ejfoundation.org
57. Anon (2002)
Death in Small Doses; Cambodia's Pesticide problems and Solutions- A report by Environmental Justice Foundation, London, UK . Internet site- www.ejfoundation.org.
58. Chowdhari D K, Nazir A, Saxena D K (2001)
Effect of Three Chlorinated Pesticides on hsromega stress gene in transgenic *Drosophila melanogaster*. *J. Biochem. Mol. Toxicol.*15(4); 173-186.
59. Rao A S, Pillala R R (July 2001)
The Concentrations of Pesticides in Sediments from Kolleru Lake in India. *Pest Management Science*; 57(7), 620-624.
60. Romeo F Quijano (2002)
Endosulfan Poisoning in Kasargod, Kerala, India - A Report on a Fact-Finding Mission, Pesticide Action Network-Asia and the Pacific, Penang, Malaysia.
61. Demeter J, Heyndrickx A, Timperman J, Lefevere M, DeBeer J (July 1977)
Toxicological Analysis in a Case of Endosulfan Suicide *Bull. Environ. Contam. Toxicol.* 18(1); 110-114.
62. Terry Gips (1990)
Breaking the Pesticide Habit- Alternatives to Twelve Hazardous Pesticides. *International Organisation of Consumers Union, Penang, Malaysia.* P 348.
63. Anon (2003)
Integrated Management of Coffee Berry Borer(Leaflet), Coffee Board, Division of Entomology/Nematology, Central Coffee Research Insitute, Coffee Research Station, Chikmangalur District, Karnataka, India.
64. Anon(2002)
Final Report of the Investigation of Unusual Illnesses allegedly produced by endosulfan exposure in Padre village of Kasaragod District (N.Kerala); National Institute of Occupational Health, Indian Council for Medical Research, Ahmedabad, India.
-

65. Vonier P M, D.DA. Crain, J A Mc Lachlan, L J Guillette Jr, Arnold S F (1996)
Interaction of Environmental chemicals with the estrogen and progesteron receptors from the oviduct of the American Alligator. Environmental Health Perstpectives 104; 1318-1322
66. Anon (Nov 12 2001)
World Wildlife Fund Comments on reregistration of Endosulfan submitted to Public Information and Record Integrity Branch, Information Resources and Services Division (7502c) Office of Pesticide Programme, Environmental Protection Agency, Washington.
67. Fransson-Steen R, Flodstram S, Warngard L (1992)
Insecticide Endosulfan and its two stereo isomers promote the growth of altered hepatic foci in rats, Carcinogenesis Vol. 13; ISS 12; P 2299-2303.
- 67a. Clarke, Harvey and Humphrey (1981)
Veterinary Toxicology (II Edition), London.
68. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Arctic Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
69. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Sub Saharan Africa- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
70. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Mediterranean- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
71. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- South East Asia and South Pacific- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
72. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Eastern and Western South America- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
73. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Central America and the Caribbean- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
74. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- North America- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
75. Anon (Dec 2002)
Regional Based Assessment of Persistent Toxic Substances- Europe- Regional Report – Chemicals- United Nations Environmental Programme- Global Environment Facility
76. Zou E, Fingerman N (1999)
Effect of Estrogenic Agents on Chitobiase activity in the epidermis and hepato pancrease of the Fiddler Crab (Uca pugilator); EcoToxicol. Environ. Saf. ; Vol.42;Iss-2; P 185-190.

76a. Zou E, Fingerman N (April 1997)

Synthetic estrogenic Agents do not Interfere with sex determination but do inhibit moulting of the Cladoceran, Daphnia magna; Bull. Environ. Contam. Toxicol.; 58(4); 594-602.

77. Cerkezkyabekir A, Aktae T (1997)

The Histopathological effects of Endosulfan on the Mouse Thyroid Gland; Turkish Journal of Biology; 21 ;P 439-444.

78. Anon (2001)

PAN Pesticide Database – Aquatic Ecotoxicity Studies for Endosulfan. (Derived from USEPA AQUIRE Acute Summaries).Pesticide Action Network, Penang, Malaysia.

79. Ciscato C H, Gebara A B, Spinosa Hde S (2002)

Pesticides residues in Cow's milk consumed in SaoPaulo City, Brazil; J. Environ. Sci. Health B.; July2002;37(4); 323-330.

80. Anon (June 2003)

Pesticide News No 60, The Journal of Pesticide Action Network UK . (Quarterly), P 19.

81. Anon (2003)

Toxicological Profile for Endosulfan ATSDR- US Dept of Health and Human Services, Public Health Services. Web site - <http://www.atsdr.cdc.gov/toxprofiles/tp41-c4.pdf>

82. Mercy Mathew, Andrews M I (Jan 2000)

Effect of Endosulfan on the hatching rate and larval survival of Common Indian Toad (Bufo melanostictus Schneider), Proceedings of the 12th Kerala Science Congress, Government of Kerala, pp 615-17

83. Anon (1999)

Global Pesticide Campaigner (Volume 09, Number 2), August 1999. (Source: Latin American News Service, April 4, 1999; [http:// www.lattn.com/cgi-local/daily.pl?094-09.](http://www.lattn.com/cgi-local/daily.pl?094-09.))

84. Devakumar C (Dec 2002)

Endosulfan Aerial Spray Controversy in Kerala. Pesticide Research Journal, Vol 14(2): pp:343-344, Society of Pesticide Science, India, Div.of Agricultural Chemicals, IARI, N.Delhi.

85. Anon (Dec 2002)

From the Editors Diary - The High Court Judgement on Endosulfan Episode of Kerala - Excerpts. Pesticide Research Journal, Vol 14(2): pp: 350-354, Society of Pesticide Science, India, Div.of Agrichem, IARI, N.Delhi.

86. Ghadiri H and Rose C W (2001)

Degradation of endosulfan in a clay soil from cotton farms of Western Queensland, Journal of Environmental Management, 62, pp155-169

87. Amaraneni.S.R., Pillale.RR(2001)

Concentrations of pesticide residues in tissues of fish from Kolleru lake in India. Environ.Toxicol.2001;16(6),550-6.

88. Leonard A.W, Hyna.R.V, Leigh.K.A, Le.J, Beckett.R (2001).

Fate and toxicity of endosulfan in Naomi River water and bottom sediments. J.Environ.Qual.2001May-June; 30(3):750-9

89. Heranadez.F, Serrano.R, Olea.N, Pitarch.E, Gaspar.JV. (2002)
Multiresidue determination of endosulfan and metabolic derivatives in human adipose tissue using automated liquid chromatographic cleanup and gas chromatographic analysis. *J. Anal. Toxicol.* 2002.Mar. 26(2):94-103.
90. Delorenzo.ME, Taylor.LA, Lund.S.A, Pennington.PL, Strozier.ED, Fulton.MH (2002)
Toxicity and bio-concentration potential of agricultural pesticide endosulfan in phytoplankton and zooplankton. *Arch. Environ. Contam. Toxicol.* 2002.Feb; 42(2):173-81.
91. Tandon.RS, L.S. Lal, Narayana.Rao.V.V,(1988)-
Interaction Endosulfan and Malathion with the green algae *Anabaena* and *Autosira fertilissima* . *Environ pollute* .1998;52(1);1-9
92. Cerejiera M J , Viana P, Batista S , Pereira T , Silva E, Valerio M J , Silva A , Ferreira M , Silva Fernandes A M (2003)
Pesticide in Portuguese surface and ground water . *Water res.* 2003 Mar; 37 (5) : 1055- 63.
93. Klemens J A, Wierland ML, Flomagin V J, Frick J A, Harper RG (2003)
A cross taxa survey of organochlorine pesticide contamination in a Costa Rica Wet land. *Environ pollute* 2003; 122(2): 245-51
94. Kutluhan.s, Akhan.G, Gulterkin.F, Kurdoglu E(2003).
Three cases of recurrent epileptic seizures caused by endosulfan. *Neurol.India.*2003. Mar: 51(1); 102-3.
- 95 . Gonzalez m, Miglitoranza K S , Aizpun De Moreno J. Emoreno V J .
Organo Chlorine Pesticide residue in leeks (*Allium porum*) Groups grown in untreated Soils from an agriculture environment. *J Agric. Food Chemi* .2003 Aug 13; 51 917) ; 5024 -9
- 96.Miglitoranza K S , de Moreno V J .
Organo chlorine pesticide sequestered in the aquatic macrophyte *Schoenoplectus ediformicus*(C.A.Meyer) from a shallow lake in Argentina. *Water Res.*2004 April; 38(7):1765-72.
- 97.Fatoki O.S., Awofolus OR (2004).
Levels of organochlorine pesticide residues in marine, surface, ground and drinking water from eastern Cape Province in South Africa. *J. Environ.Sci.Health*; 2004 Jan:39(1);101-114
98. RobertsDM., Karunarathna.A, Buckley.NA., Manuweera.G, Sheriff.MH, Eddleston.M (2003)
Influence of pesticide regulation on acute poisoning deaths in Srilanka. *Bull. World Health Organ.* 2003. 81(11):789-98.
99. Swartz.CP, Donnelly.KC, IslamZadeh.A, Rowe.GT, Rogers.WJ, Paletnikov.GM, Mekhtiev.A.A, Kasimov.R, McDonald.TJ, Wickcliffe J K , PresleyBJ BickhamJW(2003)
Chemical Contaminants and their Effects in fish and wildlife from the industrial zone of Sumgayit, Republic of Azerbaijan, *Ecotoxicology* 2003 Dec;12(6);509-21
100. Mor F, Ozmen O (2003)
Acute endosulfan poisoning in cattle. *Vet.Hum.Toxicol.* 2003 Dec; 45(6):323-4

101. Kelch JW, Kerr LA (1997)

Acute toxicosis in cattle sprayed with endosulfan. *Vet.Hum.Toxicol* 1997 Feb;39(1);29-30

102. Oktay C, Gokten E, Bozdmir N, Soyuncus(2003)

Unintentional toxicity due to endosulfan :a case report of two patients and characteristics of endosulfan toxicity . *Vet.Hum.Toxicol* 2003 Dec;45(6);318-20.

103. Rohr JR, Elskus AA, Shepherd BB, Crowley PH, Mc Carthy TM, Niedzwiecki JH, Sager T, Sih A, Palmer BD(2003)

Lethal and sublethal effects of adiazine, carbaryl, endosulfan and octylphenol on the streamside salamander (*Ambystoma barbouri*) *Environ Toxicol Chem* 2003 Oct;22(10);2385-92.

104. Hoh E, Hite RA (2004)

Sources of toxaphene and other organochlorine pesticides in North America as determined by air measurements and potential source contribution function analysis. *Environ Sci Technol* 2004 Aug 1;38(15):4187-94.

105. Carvalho FP, Gonzalez Faria F, Villeneuve JP, Catilini C, Hernandez -Garza M, Mee LD, Fowler SW (2002)

Distribution, fate and effects of pesticide residues in tropical coastal lagoons of north western Mexico. *Environ Technol.* 2002. Nov;23(11):1257-70

106. Dewan A, Bhatnagar VK, Mathur ML, Chakma T, Kashyap R, Sandhu HG, Sinha SN, Saiyed HN (2004)

Repeated Episodes of Endosulfan poisoning. *J. Toxicol.Chin Toxicol.* 2004;4294): 363-9

107. Grumfeld HT, Bonefeld-Jorgensen EC (2004)

Effects of invitro estrogenic pesticides on human oestrogen receptor alpha and beta mRNA levels. *Toxicol.Lett* 2004 Aug 1;151(3);467-80.

108. Kalendar S, Kalendar Y, Ogutcu A, Uzunhisarcikli M, Durak D, Acikgoz F (2004)

Endosulfan-induced cardio-toxicity and free radical metabolism in rats.the protective effect of Vitamin E. *Toxicology.* 2004 Oct 1;202(3);227-35.

109. Sutherland TD, Home I, Weir KM, Russel RJ, Oakeshott JG (2004),

Toxicity and residues of endosulfan isomers. *Rev.Environ.Cotam.Toxicol;* 2004;183:99-113.

110. Wade MG, Desaulniers D, Leingartnerk, Foster WG(1997)

Interactions between endosulfan and dieldrin on estrogen mediated processes invitro and invivo. *Reprod.Toxicol.* 1997 Nov-Dec;11(6);791-8.

111. Garg UK, Pal AK, Jha GJ, Jadhao SB (2004)

Haemato-biochemical and immunopathophysiological effects of chronic toxicity with synthetic pyrethroid, organophosphate and chlorinated pesticides in broiler chicks. *Int.Immuno Pharmacol.* 2004 Dec 4 (13);1709-22

112. Prados Rosales RC, Luque Garcia JL, Luque de Castro MD

Rapid Analytical method for the determination of pesticide residues in sunflower seeds based on focussed microwave-assisted Soxhlet extraction prior to gas chromatography-tandem mass spectrometry. *J.Chromatogr.A.* 2003 Apr 18;993(1-2):121-9

113. Miglioranza KS, Aizpun de Moreno JE, Moreno VJ
Land based sources of marine pollution: organochlorine pesticide in stream system. *Environ Sci Pollut Res Int* 2004;11(4);227-32.
114. Goulet BN, Hastela A (2003)
Toxicity of cadmium, endosulfan and atrazine in adrenal steroidogenic cells of two amphibian species, *Xenopus laevis* and *Rana catesbeiana* *Environ Toxicol Chem* 2003 Sep;22(9)2;106-13
115. Vorkamp K, Riget F, Glasius M, Pecseli M, Lebeuf M, Muir D. (2004)
Chlorobenzene, chlorinated pesticide coplanar chlorobiphenyl, and other organochlorine compounds in Greenland Gota. *Sci Total Environ* 2004 Sep 20;331(1-3);1
116. Chiasson H, Vincent C, Bostanian NJ
Insecticide properties of a chenopodium based botanical. *J Econ Entomol.* 2004. Aug;97(4) :1378-83.
117. Ibarluzea Jon J, Fernandez MF, Santa Marina L, Olea Serrano MF, Rivas AM, Aurrekoetxea JJ, Enposito J, Lorenzo M, Torne P, Villalobos M, Pedraza V, Sasco AJ, Olea N (2004).
Breast cancer risk and the combined effect of environmental estrogens- *Cancer causes control.* 2004 Aug; 15 (6): 591-600
118. Habibullah Saiyed, Aruna Dewan, Vijay Bhatnagar, Udyavar Shenoy, Rathika Shenoy, Hirehall Rajmohan, Kumud Patel, Rekha Kashyap, Pradip Kulkarni, Bagalur Rajan, Bhadabhai Lakkad (2003)
Effect of Endosulfan on Male Reproductive Development, *Environmental Health Perspectives* Dec 2003 Vol 111 no.16.
119. Jamil K, Shaik AP, Mahboob M, Krishna D (2004)
Effect of organophosphorus and organochlorine pesticides (monochrotophos, chlorpyrifos, dimethoate and endosulfan) on human lymphocytes in-vitro. *Drug Chem Toxicol.* 2004 May;27 (2) :133-44.
120. Park EK, Lees EM (2004)
The interaction of endosulfan with the collembolan *Proistoma minuta* (Tullberg): toxicity, the effects of sublethal concentrations and metabolism. *Pest manag. Sci* .2004 Jul ; 60(7) : 710-8
121. Downing HF, De Lorenzo ME, Fulton MH, Scott GI, Madden CJ, Kucklick JR (2004)
Effects of the agricultural pesticides atrazine, chlorothalonil and endosulfan on South florida microbial assemblages. *Ecotoxicology* 2004 Apr ; 13 (3) : 245-60.
122. Jergentz S, Mugin H, Bonetto C, Schulz R (2004)
Runoff related endosulfan contaminations and aquatic macro invertebrate responses in rural basins near Buenos Aires, Argentina. *Arch. Environ Contam. Toxicol.* 2004 Apr ;46 (3) :345-352.
123. Thiere G, Schuz R (2004)
Runoff related agricultural impact in relation to macro invertebrate communities of the Lourens river, South Africa. *Water Res.* 2004 ; 38(13) 3092-102

124. Tariq M I, Afzal S, Hussian L (2004)

Pesticide in shallow ground water of Bahawal nagar, Mussafargarh, D G Khan and Rajanpur districts of Punjab, Pakistan. *Environ. Int.* 2004 June ; 30(4) ; 471-9.

125. Otludil B, Cengiz EI, Yildirim M Z, Unver O, Unlu E

The effects of endosulfan on the great ramshorn snail *Planorbis corneus* (gastropoda, pulmonata) a histopathological study. *Chemosphere* 2004. Aug; 56 (7) : 707-16.

126. Tripathi G, Verma P

Endosulfan mediated biochemical changes in the fresh water fish *Clarias batrachus*. *Biomed Environ Sci* 2004 Mar 17 (1) 47-56.

127. Zhang Z, Huang J, Yu G, Hong H (2004)

Occurrence of PAHs, PCBs, and organochlorine pesticides in the Tonghui River of Beijing, China. *Environ Pollut* 2004. Jul ; 130 (2) ; 249-61.

128. Sanghi R, Pillai MK, Jayalakshmi TR, Nair A

Organochlorine and organo-phosphorous pesticide residues in breast milk from Bhopal, Madhya Pradesh, India. *Human. Exp. Toxicol.* 2003 Feb ; 22 (2) 73-6.

129. Mosleh YY, Paris-Palacois S, Couderchet M, Vernet G. (2003)

Acute and sub-lethal effects of two insecticides on earthworm (*Lumbricus terrestris* L) under laboratory conditions. *Environ Toxicol.* 2003 Feb ; 18 (1) ; 1-8.

130. Gonzalez-Farias F, Gisneros Estrada X, Fuentes Rui Z C, Diaz Gonzalez G, Botello AV (2002)

Pesticides distribution in sediments of tropical coastal lagoon adjacent to an irrigation district in northwest Mexico. *Environ. Technol.* 2002 Nov ; 23 (11); 1247-56

131. Gamon M, Saez E, Gil J, Boluda R (2003)

Direct and indirect exogenous contamination by pesticides of rice farming soils in a Mediterranean wetland. *Arch. Environ. Contam. Toxicol.* 2003 Feb ; 44 (2) ; 141-51.

132. Michael Hermann (2003)

Endosulfan Preliminary Dossier ; www.unece.org/env/popsxg/docs/2000-2003/dossier-endosulfan-may03.pdf.

133. Medina, JR et al. (1991)

Insecticide residues in selected well water in Calamba and Calauan, Laguna. *The Philippine Agriculturist* 74(2):195-205, 1991.

134. Paningbatan, EP et al. (1993).

Movement of monocrotophos and endosulfan in saturated soil. *The Philippine Agriculturist* 76(3): 259-268.

<http://www.ipen.org>